

Nomenclature of Proteins of Cow's Milk: Fifth Revision¹

W. N. EIGEL, Chairman

Department of Food Science and Technology
Virginia Polytechnic Institute and State University
Blacksburg 24061

J. E. BUTLER

Department of Microbiology
College of Medicine
University of Iowa
Iowa City 52240

C. A. ERNSTROM

Department of Food Science
Utah State University
Logan 84321

H. M. FARRELL, JR.

Eastern Regional Research Center
Agricultural Research Service, US Department of Agriculture
Philadelphia 19118

V. R. HARWALKAR

Food Research Institute
Agriculture Canada
Ottawa, Canada

R. JENNESS

Department of Biochemistry
University of Minnesota
St. Paul 55108

R. McL. WHITNEY

Department of Food Science
University of Illinois
Urbana 61801

ABSTRACT

This report reviews changes the nomenclature of bovine milk proteins necessitated by recent advances of our knowledge. Identification of a number of milk proteins (α_{S1} -, β -, and κ -caseins; α -lactalbumin and β -lactoglobulin) continues to be based upon their primary structures (amino acid sequences). Since our last report, α_{S2} -casein and serum albumin can be added to the list of major milk proteins

for primary structure is known. Changes recommended in the nomenclature of caseins are primarily a result of differences within this family of proteins brought about by posttranslational modification. For example, α_{S0} -casein is identical to α_{S1} -casein, and α_{S3} -, α_{S4} -, and α_{S6} -caseins are identical to α_{S2} -casein except for differences in degree of phosphorylation. Additionally, proteose-peptone components 5, 8-slow and 8-fast, and γ_1 -, γ_2 -, and γ_3 -caseins are N-terminal and C-terminal fragments, respectively, of β -casein formed during proteolysis by plasmin. Nomenclature of immunoglobulins remains consistent with guidelines for human proteins and is based

Received August 22, 1983.

¹Report of the Committee on the Nomenclature and Methodology of Milk Proteins of the Dairy Foods Research Section, Dairy Foods Division, American Dairy Science Association.

TABLE 1. Proteins of bovine milk and some of their properties.*

Protein and suggested abbreviation	Composition in skim milk (g/liter)	Genetic variants	Molecular ^c weights	Sedimentation ^d constant	Isoionic ^e point	Isoelectric ^e point	A 1% ^f 1 cm	H ₂ O ^g avg. (kcal/residue)
α_{s1} -Casein (α_{s1} -CN)	12-15	A	22,068	...	5.15 (243) ^h	...	10.10 (244)	1056
		B	23,614	1.64 (225)	4.92-5.05 (33,233,243)	4.44-4.76 (252)	10.05 (244)	1061
		C	23,542	1.64 (225)	5.00-5.35 (233,243)	...	10.0 (258)	1061
		D	23,724	10.03 (244)	1060
		E	23,542	...	5.16 ^h (119)	4.2-4.6 ^h (138)	10.7 ⁱ , 10.0 ^h (9,194)	1068
α_{s2} -Casein (α_{s2} -CN)	3-4	A	25,230 ^j	997
		B
		C
		D
β -Casein (β -CN)	9-11 ^b	A ¹	24,023	1.51, 1.50 (77,195)	5.41 (233)	...	4.6, 4.7 (245,258)	1221
		A ²	23,983	1.51, 1.50 (77,195)	5.30 (233)	4.83-5.07 (252)	...	1231
		A ³	23,974	...	5.20 (233)	1229
		B	24,092	...	5.53 (233)	...	4.7 (245)	1223
		C	23,944	...	5.85 (233)	...	4.5 (245)	1228
		D	23,944	1240
		E	23,982	...	5.35 ^h (119)	4.6-5.1 ^h (138)	...	1238
κ -Casein (κ -CN)	2-4	A	19,039	...	5.77 (233)	5.45-5.77 (252)	...	1094
		B	19,007	1.4 ^h (237)	6.07 (233) 5.37 ^h (119)	4.1 ^h (235) 5.3-5.8 ^h (138)	10.5 ^{h,i} (91)	1112
β -Lactoglobulin (β -LG)	2-4	A	18,363	...	5.35 (180)	5.13 (88)	9.6 (248)	1080
		B	18,277	...	5.41 (180)	5.13 (88)	10.0, 9.6 ⁱ (135,248)	1074
		C	18,286	...	5.39 (180)	...	9.5 (248)	1077
		D	18,276	1077
		E	18,205	1074
		F	18,243
		G	18,223	8.7 ⁱ (24)	...
		(Dr)	...	2.7, 1.9 ^h (248,249,250)	...	5.14 ^h (213)	10.8 ^h (27)	...
α -Lactalbumin (α -La)	.6-1.7	A	14,147	1016
		B	14,175	1.92 ^h , 1.98 ^h , 1.87 ^h (156,240,262)	...	4.2-4.5 ^h (156,157)	20.9 ^h , 20.1 ^h , 20.6, 20.1 ^h (69,156,211,227)	1022
Serum Albumin (SA)	.4	A	66,267	5.01, 4.0 (15,75)	5.13 (119) 5.15 (222)	4.7-4.9 (213)	6.9, 6.6, 6.49, 6.3 (140,203,215,260)	995

Immunoglobulin G ₁ (IgG ₁)	.3-.6	153,000-163,000 (28,68,148,164,241)	6.3-7.0 (49,164,241)	...	5.5-6.8 (139)	12.1 (241) 13.5 (164)	...
Immunoglobulin G ₂ (IgG ₂)	.05-.1	146,000-154,000 (68,148,164,241)	6.5-7.1 (49,164,241)	...	7.5-8.3 (139)	12.1 (241) 12.0 (164)	...
Immunoglobulin A ^k (IgA)	.05-.15	385,000-417,000 (50,68,169)	10.8-11.0 (50,68,168,205)
Immunoglobulin M (IgM)	.05-.1	1,000,000 (188) 960,000 (143)	18.2-19.8 (143,158,168,188)	11.8 (241)	...
Secretory Component (SC)	.02-.1	79,000	4.0 (50) 4.9 (168)

^aInclusion of values in this Table does not constitute endorsement by the committee.

^bValues given for β - and γ -caseins were combined and reported as β -CN in this table.

^cAll values, except for immunoglobulins, were calculated based upon known amino acid composition.

^dSedimentation constant, S_{20W} , in Svedberg units (1×10^{-13} sec).

^eSee original references for details of methodology.

^fAbsorptivity of a 1% solution measured in a 1 cm light path at 280 nm, except where noted otherwise.

^gAverage hydrophobicity, calculated using free energies of transfer of amino acid side chains from an organic environment to an aqueous environment (30) and known amino acid compositions of the various proteins.

^hGenetic variant of protein was not reported.

ⁱAbsorptivity measured at 278 instead of 280 nm.

^jMolecular weight calculated for α_{S2} -CN 11 P.

^kAll physical chemical data obtained with secretory IgA.

largely upon crossreactivity with reference proteins.

The minor whey protein lactollin is β_2 -microglobulin for which the sequence of amino acids is known. An operational definition for proteins associated with the milk fat globule membrane has been developed. Nomenclature initially suggested for these proteins was based upon their electrophoretic behavior under a given set of conditions.

Because of increased interest in milk proteins of species other than bovine, the Committee suggests that these be identified as homologs of those already characterized in European, *Bos taurus*, and Indian, *Bos indicus*, cattle. Guidelines are given to aid in determining if homology exists. Provisional nomenclature is suggested for use in the interim until homology can be established.

INTRODUCTION

The initial report of the American Dairy Science Association Committee on the Nomenclature, Classification, and Methodology of Milk Proteins (129) was an attempt to clarify the nomenclature of milk proteins by "presenting a summary of preferred usage and by showing the relationship between the individual proteins which have been isolated and the classical fractions." Subsequently, this Committee has published a revision of milk protein nomenclature approximately every 5 yr (48, 218, 246, 267) to summarize more recent findings (Table 1) and to suggest changes in nomenclature where appropriate. The intent of this Committee is to suggest a flexible nomenclature system that allows for incorporation of new discoveries rather than to suggest prematurely a rigid system of nomenclature. Since the last report of this Committee (267), the most significant findings related to nomenclature are: determination of primary sequences of α_{s2} -casein and serum albumin, demonstration that members of the α_{s1} - and α_{s2} -casein families (and possibly κ -casein) differ in degree of phosphorylation, determination of the source of several minor proteins in milk as fragments of β -casein produced by plasmin, and identification of lactollin as β_2 -microglobulin and determination of its primary sequence. Sug-

gested changes in the nomenclature of the major proteins allow for differentiation based upon differences resulting from posttranslational modifications.

For the first time this Committee recommends an operational nomenclature for proteins associated with the milk fat globule membrane as well as a system, based upon electrophoretic behavior, for identifying individual components. Although many of the proteins associated with the milk fat globule membrane possess enzymatic activity (4, 45, 196), we have elected not to list them, which is in keeping with past practices of not elaborating on enzymes associated with skim milk. These enzymes have been reviewed elsewhere (228).

With the exception of lactollin, we have made no mention of a number of minor milk proteins reviewed by Groves (106) and have followed the pattern previously established (48, 129, 218, 246, 265) in limiting the scope of this report. More detailed treatments on specific aspects of the milk proteins are available in several recent reviews (46, 47, 55, 83, 84, 163, 185, 226, 266). The present report breaks tradition with previous ones by suggesting nomenclature for milk proteins isolated from species other than *Bos*. Because of the increased interest in milk proteins of other species and the tendency of researchers to name these proteins after their counterparts in *Bos*, we felt a responsibility to establish criteria for naming these proteins and determining their homology with bovine milk proteins. At the same time, we respect the prerogative of any investigator to assign nomenclature to any protein with no counterpart in the milks of European, *Bos taurus*, and Indian, *Bos indicus*, cattle.

THE CASEINS

Caseins in milk of the genus *Bos* were defined originally by this Committee (129) as those phosphoproteins that precipitate from raw skim milk by acidification to pH 4.6 at 20°C. In a subsequent report (267), the Committee differentiated caseins according to their relative electrophoretic mobility in alkaline polyacrylamide or starch gels containing urea with or without mercaptoethanol. Based upon current knowledge, we now recommend that use of electrophoresis as a basis for classification be dropped and that caseins be identified

according to homology of their primary structures (amino acid sequences) into the following families: α_{s1} -, α_{s2} -, β -, and κ -caseins. Individual members of these families still can be identified by gel electrophoretic techniques, some of the more effective of which are suggested in the monograph by this Committee (238).

α_{s1} -Caseins

The α_{s1} -caseins (α_{s1} -CN) consist of one major and one minor component, both with the same amino acid sequence established by Mercier et al. (187) and Grosclaude et al. (103) for the major component. At present, the five genetic variants are designated A, D, B, C, and E in order of decreasing relative electrophoretic mobilities in alkaline gels containing urea (100, 199, 242). The B variant is predominant in *Bos taurus*, and C is in *Bos indicus* and *Bos grunniens* (99, 242). Bell et al. (23) observed an electrophoretic band in patterns of skim milk protein from Bali (Banteng) cattle, *Bos javanicus*, with slower mobility than the C variant. However, more information is needed before this band can be designated as a unique genetic variant of α_{s1} -CN.

The minor component, previously classified as α_{s0} -casein (267), has the same amino

acid sequence as α_{s1} -CN but contains one additional phosphorylated serine residue at position 41 (174, 175). We recommend that variations, which result from posttranslational modification, such as phosphorylation, be designated by an Arabic number and a letter after the Latin letter that designates the genetic variant. In this case, the letter P will indicate that the posttranslational variation is a result of differences in amounts of phosphorylation (31, 32, 100), and the Arabic number will indicate the number of phosphates attached. Thus, the major genetic variant in *Bos taurus* would be designated α_{s1} -CN B-8P instead of α_{s1} -CN B, and the term α_{s0} -casein B would be dropped in favor of α_{s1} -CN B-9P. Because the two α_{s1} -CN components differ only in degree of phosphorylation, genetic variants of both components will be identical in a specific milk. When these proteins are prepared from milk of heterozygous animals in which the particular casein genes show no dominance, two genetic variants will be present.

The primary structure of α_{s1} -CN B-8P is in Figure 1. It consists of 199 amino acid residues: Asp₇, Asn₈, Thr₅, Ser₈, Ser P₈, Glu₂₄, Gln₁₅, Pro₁₇, Gly₉, Ala₉, Val₁₁, Met₅, Ile₁₁, Leu₁₇, Tyr₁₀, Phe₈, Lys₁₄, His₅, Trp₂, and Arg₆ with

H.Arg - Pro - Lys - His - Pro - Ile - Lys - His - Gln - Gly -										10
Leu - Pro - Gln -										14
Glu - Val - Leu - Asn - Glu - Asn - Leu -										20
										Absent in Variant A
Leu - Arg - Phe - Phe - Val - Ala -										26
Pro - Phe - Pro - Gln - Val - Phe - Gly - Lys - Glu - Lys - Val - Asn - Glu - Leu -										30
Ser - Lys - Asp - Ile - Gly - Ser - Glu - Ser - Thr - Glu - Asp - Gln -										41
P (Variant B-9P)										53
Ala - Met - Glu - Asp - Ile - Lys -										59
Gln - Met -										60
Glu - Ala - Glu - Ser - Ile - Ser - Ser - Ser - Glu - Glu - Ile - Val - Pro - Asn - Ser - Val - Glu - Gln - Lys - His -										70
P										80
Ile - Gln - Lys - Glu - Asp - Val - Pro - Ser - Glu - Arg - Tyr - Leu - Gly - Tyr - Leu - Glu - Gln - Leu - Leu - Arg -										90
Leu - Lys - Lys - Tyr - Lys - Val - Pro - Gln - Leu - Glu - Ile - Val - Pro - Asn - Ser - Ala - Glu - Glu - Arg - Leu -										110
P										120
His - Ser - Met - Lys - Glu - Gly - Ile - His - Ala - Gln - Gln - Lys - Glu - Pro - Met - Ile - Gly - Val - Asn - Gln -										130
Glu - Leu - Ala - Tyr - Phe - Tyr - Pro - Glu - Leu - Phe - Arg - Gln - Phe - Tyr - Gln - Leu - Asp - Ala - Tyr - Pro -										150
Ser - Gly - Ala - Trp - Tyr - Tyr - Val - Pro - Leu - Gly - Thr - Gln - Tyr - Thr - Asp - Ala - Pro - Ser - Phe - Ser -										170
Asp - Ile - Pro - Asn - Pro - Ile - Gly - Ser - Glu - Asn - Ser -										190
Glu - Lys - Thr - Thr - Met - Pro - Leu - Trp.OH										192
Gly (Variants C & E)										199

Figure 1. Primary structure of *Bos* α_{s1} -CN B-8P (103, 187). The enclosed amino acid residues are sites corresponding to mutational differences in the genetic variants: A, C, D, and E (100, 102). The enclosed P represents the site of additional phosphorylation in α_{s1} -CN B-9P.

a calculated molecular weight of 23,614 (103, 187). The other genetic variants differ from B, as indicated in Figure 1 and Table 2. For variant D, the appropriate nomenclature should be α_{s1} -CN D-9P for the major component and α_{s1} -CN D-10P for the minor (Figure 1, Table 2).

Peptides released during proteolysis of α_{s1} -CN by plasmin have identical gel electrophoretic mobilities under alkaline conditions in the presence of urea as peptides extracted from casein with N,N-dimethylformamide (3). These extracted peptides have complex electro-

phoretic patterns identical with λ -casein (76), an ill-defined fraction first prepared by Long et al. (165). Two of the peptides produced by plasmin degradation of α_{s1} -CN had tryptic peptide maps and molecular weights identical with two peptides from the N,N-dimethylformamide extract with corresponding electrophoretic mobilities. Although the fraction once identified as λ -casein by this Committee (246) now appears to consist of fragments of α_{s1} -CN, much more definitive work is needed before additional changes are made in the nomenclature.

TABLE 2. Location of amino acid substitutions in genetic variants of milk proteins identified in *Bos*.

TABLE 2. Location of amino acid substitutions in genetic variants of human α_1 -antitrypsin										
Protein	Variant	Position								
		14-26		53		59		192		
α_{s1} -CN	A	Deleted		Ala		Gln		Glu Gly		
	B									
	C									
	D									
	E									
β -CN	A ¹	18	35	36	37	67	106	122		
	A ²	SerP	SerP	Glu	Glu	His Pro	His Gln	Ser		
	A ³					His		Arg		
	B		Ser		Lys	His				
	C									
	D	Lys								
	E			Lys						
					136	148				
κ -CN	A					Thr				
	B					Ile	Asp Ala			
α -LA	A					10				
	B					Gln Arg				
β -LG		45	50	59	64	78	118	129	130	158
	A				Asp		Val			
	B	Glu	Pro	Gln	Gly	Ile	Ala	Asp	Asp	Glu
	C			His						
	D	Gln								
	E (Dyak)									Gly (Gly) ^a
	F		(Ser) ^a					(Tyr) ^b	(Tyr) ^b	(Gly) ^a
	G					(Met) ^a				(Gly) ^a

^aAmino acid substitutions placed in parentheses have not yet been established by sequencing the genetic variant.

^bTyr substitution for Asp occurs at either 129 or 130.

10
 H. Lys - Asn - Thr - Met - Glu - His - Val - Ser - Ser - Ser - Glu - Glu - Ser - Ile - Ile - Ser - Gln - Glu - Thr - Tyr -
 20
 Lys - Gln - Glu - Lys - Asn - Met - Ala - Ile - Asn - Pro - Ser - Lys - Glu - Asn - Leu - Cys - Ser - Thr - Phe - Cys -
 30
 Lys - Glu - Val - Val - Arg - Asn - Ala - Asn - Glu - Glu - Glu - Glu - Tyr - Ser - Ile - Gly - Ser - Ser - Ser - Glu - Glu -
 40
 Ser - Ala - Glu - Val - Ala - Thr - Glu - Glu - Val - Lys - Ile - Thr - Val - Asp - Asp - Lys - His - Tyr - Gln - Lys -
 50
 Ala - Leu - Asn - Glu - Ile - Asn - Glu - Phe - Tyr - Gln - Lys - Phe - Pro - Gln - Tyr - Leu - Gln - Tyr - Leu - Tyr -
 60
 Gln - Gly - Pro - Ile - Val - Leu - Asn - Pro - Trp - Asp - Gln - Val - Lys - Arg - Asn - Ala - Val - Pro - Ile - Thr -
 70
 Pro - Thr - Leu - Asn - Arg - Glu - Gln - Leu - Ser - Thr - Ser - Glu - Glu - Asn - Ser - Lys - Lys - Thr - Val - Asp -
 80
 Met - Glu - Ser - Thr - Glu - Val - Phe - Thr - Lys - Lys - Thr - Lys - Leu - Thr - Glu - Glu - Glu - Lys - Asn - Arg -
 90
 Leu - Asn - Phe - Leu - Lys - Lys - Ile - Ser - Gln - Arg - Tyr - Gln - Lys - Phe - Ala - Leu - Pro - Gln - Tyr - Leu -
 100
 Lys - Thr - Val - Tyr - Gln - His - Gln - Lys - Ala - Met - Lys - Pro - Trp - Ile - Gln - Pro - Lys - Thr - Lys - Val -
 110
 Ile - Pro - Tyr - Val - Arg - Tyr - Leu. OH
 207

Figure 2. Primary structure (41, 42) of *Bos* α_{s2} -CN A-11P. The amino acid sequences enclosed in brackets represent the possible amino acid sequence missing in the D variant. Only 11 of the phosphorylated serine residues have been tentatively identified (42) and have been designated with a P.

α_{s2} -Caseins

Since the last report of this Committee (267) the primary structure of α_{s2} -casein (α_{s2} -CN) (Figure 2) has been determined (41, 42). It consists of 207 amino acid residues: Asp₄, Asn₁₄, Thr₁₅, Ser₆, Ser P₁₁, Glu₂₅, Gln₁₅, Pro₁₀, Gly₂, Ala₈, Cys₂, Val₁₄, Met₄, Ile₁₁, Leu₁₃, Tyr₁₂, Phe₆, Lys₂₄, His₃, Trp₂, and Arg₆ with a calculated molecular weight of 25,230 (42, 97). Other milk protein components, previously classified as α_{s3} -, α_{s4} -, and α_{s5} -casein by this Committee (218, 267) and α_{s6} -casein by Annan and Manson (9), appear to be components of the α_{s2} -CN family. Evidence strongly suggests that all have the same amino-acid sequence but differ in phosphate content (41, 42). The exact location of the phosphate moieties in each member of the family of α_{s2} -CN has not been determined. Those positions shown as phosphorylated (Figure 2) occur in at least one of the α_{s2} -CN fractions (42). We, therefore, tentatively recommend, until further information is available as to their degree of phosphorylation, that the nomenclature of the A variant, for example, be changed as shown in Table 3. The α_{s5} -casein is a dimer consisting of α_{s3} - and α_{s4} -caseins, now called α_{s2} -CN A-12P and α_{s2} -CN A-11P, linked together by a disulfide bond (120). Until

location of this disulfide bond is shown to be unique to these particular fractions, we recommend deletion of this term.

The four recognized genetic variants of α_{s2} -CN are designated A, B, C, and D. The A and D variants have been observed in European breeds (*Bos taurus*) with D in the Vosgienne and Montbéliarde breeds (97). The electrophoretic mobilities of the various D variant bands are slower than the corresponding bands in the A variant at pH 8.6 but are faster at pH 3.0 (97, 98). In addition to variant A, variant B is found in *Bos taurus* and *Bos indicus* in a high Nepalese valley, and variant C is observed specifically in yaks (*Bos grunniens*) in the same region (101). The C variant was also found in yaks from

TABLE 3. Summary of changes recommended for the nomenclature of the α_{s2} -CN A family.

Former nomenclature	Recommended nomenclature
α_{s2} -CN A	α_{s2} -CN A-13P
α_{s3} -CN	α_{s2} -CN A-12P
α_{s4} -CN	α_{s2} -CN A-11P
α_{s6} -CN	α_{s2} -CN A-10P

the Republic of Mongolia and the B variant in zebu from the Republic of South Africa (98).

Differences among primary sequences of polypeptide chains in genetic variants of α_{S2} -CN, and exact locations and, perhaps, even number of phosphate groups in the various components still remain to be established. Grosclaude et al. (101) compared amino acid and phosphate contents of the C variant with α_{S2} -CN A-10P (Table 4) and postulate a possible substitution of glycine for a phosphorylated serine. Variant D, like α_{S1} -casein A, has a peptide deleted (97, 98); nine amino acid residues are missing from the sequence of α_{S2} -CN A-11P. The exact peptide has not been established because of similarities of the three possibilities: from residues 50 to 58 inclusive, from residues 51 to 59, or from residues 52 to 60 (Figure 2). Because this peptide contains three phosphorylated serine residues, the D variant component corresponding to a α_{S2} -CN A-11P should be named α_{S2} -CN D-8P.

β -Caseins

The primary structure of β -casein (β -CN) (Figure 3) has been elucidated by Ribadeau-Dumas and coworkers (103, 216). Based on this structure, Groves et al. (95, 109, 110) showed that those proteins previously called γ_1 -, γ_2 -, and γ_3 -caseins (267) are actually fragments of β -CN consisting of residues 29-209, 106-209, and 108-209, respectively. Plasmin, a protease in milk (73, 141) is responsible for proteolysis of β -CN, leading to formation of γ_1 -, γ_2 -, and γ_3 -caseins (71). The N-terminal fragments of β -CN released during plasmin degradation also have been identified in milk. Andrews (7, 8) showed that proteins previously called proteose peptone components 5 and 8-fast (267) are identical with fragments of β -CN. Proteose peptone 8-fast consists of residues 1-28, and component 5 is actually the two fragments comprising residues 1-105 and 1-107. Fragments released during proteolysis of

TABLE 4. Comparison of the amino acid and phosphate content of comparable components^a of variant A and C of α_{S2} -CN (101).

Amino acid	Variants	
	A (residues/molecule)	C (residues/molecule)
Aspartic acid	18	18
Threonine	15	15
Serine	17	11 ^{b,c}
Proline	10	13 ^c
Glutamic acid	40	40
Glycine	2	3 ^c
Alanine	8	8
Valine	14	14
Methionine	4	4
Isoleucine	11	11
Leucine	13	14 ^c
Tyrosine	12	12
Phenylalanine	6	6
Tryptophane	2	...
Lysine	24	24
Histidine	3	3
Arginine	6	6
Cystine	2	...
H ₂ PO ₄	10	8 ^c

^aComponent previously designated α_{S6} -CN.

^bLow value may be an artifact of acid hydrolysis.

^cDifferent in amino acid or phosphate content from A variant.

β -CN with plasmin are identical to proteose peptone component 5 (72). The remaining fragments of β -CN (residues 29-105 and 29-107) released during plasmin degradation are proteose peptone component 8-slow (74). Whether proteolysis of β -CN by plasmin occurs primarily in mammary epithelial cells, during temporary storage in the alveolar and glandular lumina, during refrigerated storage, or during protein isolation and purification is not known. Nevertheless, on the basis of these advances in our knowledge, we recommend that fragments resulting from proteolytic cleavage be named as

derivatives of the parent polypeptide from which they were derived. A summary of the new recommended nomenclature for the β -CN family is in Table 5.

Seven genetic variants of β -CN are known, but their differentiation by gel electrophoresis is more complicated than for the other caseins. In alkaline gels containing urea (149), they migrate in the order $A^1=A^2=A^3>B>D,E>C$. However, under acidic conditions (149), their order is $C>B=D>A^1=E>A^2>A^3$. Thus, although A variants can be differentiated from B, C, D, and E by electrophoresis under alkaline con-

TABLE 5. Changes recommended in nomenclature of the β -CN family.

Former nomenclature (267)	Recommended nomenclature	Source of fragment
β -Casein A ¹	β -CN A ¹ -5P	...
β -Casein A ²	β -CN A ² -5P	...
β -Casein A ³	β -CN A ³ -5P	...
β -Casein B	β -CN B-5P	...
β -Casein C	β -CN C-4P	...
β -Casein D	β -CN D-4P	...
β -Casein E	β -CN E-5P	...
γ_1 -Casein A ¹	β -CN A ¹ -1P (f29-209)	β -CN A ¹ -5P
γ_1 -Casein A ²	β -CN A ² -1P (f29-209)	β -CN A ² -5P
γ_1 -Casein A ³	β -CN A ³ -1P (f29-209)	β -CN A ³ -5P
γ_1 -Casein B	β -CN-B-1P (f29-209)	β -CN B-5P
γ_2 -Casein A ²	β -CN A ² (f106-209)	β -CN A ¹ -5P or β -CN A ² -5P
γ_2 -Casein A ³	β -CN A ³ (f106-209)	β -CN A ³ -5P
γ_2 -Casein B	β -CN B (f106-209)	β -CN B-5P
γ_3 -Casein A	β -CN A (f108-209)	β -CN A ¹ -5P, β -CN A ² -5P or β -CN A ³ -5P
γ_3 -Casein B	β -CN B (f108-209)	β -CN B
Proteose peptone ^a component 5	β -CN-5P (f1-105)	β -CN
Proteose peptone ^a component 5	β -CN-5P (f1-107)	β -CN
Proteose peptone ^a component 8-slow	β -CN-1P (f29-105)	β -CN
Proteose peptone ^a component 8-slow	β -CN-1P (f29-107)	β -CN
Proteose peptone ^a component 8-fast	β -CN-4P (f1-28)	β -CN

^aThese fragments have not been isolated from specific genetic variants. When their genetic origins and degree of phosphorylation are established, appropriate letters and numerals should be employed.

ditions, acidic conditions are required for differentiation of A variants. An additional electrophoretic band with mobility slower than β -CN A³ under acidic conditions has been observed in the milk of Bali (Banteng) cattle, *Bos (Bibos) javanicus* (23). Although this electrophoretic band has been called β -CN A⁴, additional evidence is needed prior to actual assignment of nomenclature. In 1968, Aschaffenburg et al. (14) observed a variant of β -CN in Indian and African zebu cattle (*Bos indicus*) that had the same electrophoretic behavior as the B variant but appeared to have different peptide maps. This Committee previously designated this variant B_Z (218). However, Grosclaude et al. (99) observed that the amino acid substitution in the A¹ variant was the same for *Bos indicus* and *Bos taurus* and that the B variant in Choa zebu cattle differed from the A¹ variant by substitution of an arginine residue for a serine residue at position 122, the same difference observed in *Bos taurus*. This raises a question of the existence of a variant B_Z. More work is needed before existence of this variant can be confirmed. The A variants are the predominant polymorphs in all species of *Bos* investigated (242).

The β -CN A²-5P consists of a single polypeptide chain containing 209 residues (Figure 3): Asp₄, Asn₅, Thr₉, Ser₁₁, Ser P₅, Glu₁₈, Gln₂₁, Pro₃₅, Gly₅, Ala₅, Val₁₉, Met₆, Ile₁₀, Leu₂₂, Tyr₄, Phe₉, Lys₁₁, His₅, Trp₁, and Arg₄ with a calculated molecular weight of 23,983. The other genetic variants differ (99, 102, 104) as in Figure 2 and Table 2. Again, both C and D variants have one less phosphate group than others.

Fragments (f 29-209), (f 106-209), and (f 108-209) have been detected and identified only in β -CN variants A¹, A², A³, and B (Table 5) (95, 109). Examination of milks containing the C variant indicates that the (f 29-209) fragment is missing. The charge reversal occurring in this variant at residue 37 may be related to this phenomenon. While it is expected that these fragments also would occur in variants D and E, they have not been reported specifically. Both β -CN A¹-5P and A²-5P yield the same fragment (f 106-209) because the point of genetic mutation occurs prior to residue 106. Therefore, this fragment is called β -CN A² (f 106-209) to provide a definitive nomenclature. Similarly, β -CN A¹-5P, A²-5P, and A³-5P yield the same (f 108-209) fragment, which by




Figure 3. Primary structure of *Bos* β -CN A²-5P (103, 216). The enclosed amino acid residues are the sites corresponding to the mutational (99, 102, 104) differences in the genetic variants: A¹, A³, B, C, D, and E. The arrows indicate the points of attack by plasmin responsible for β -CN fragments present in milk.

10
 PyroGlu - Glu - Gln - Asn - Gln - Glu - Gln - Pro - Ile - Arg - Cys - Glu - Lys - Asp - Glu - Arg - Phe - Phe - Ser - Asp -
 20
 Lys - Ile - Ala - Lys - Tyr - Ile - Pro - Ile - Gln - Tyr - Val - Leu - Ser - Arg - Tyr - Pro - Ser - Tyr - Gly - Leu -
 30
 Asn - Tyr - Tyr - Gln - Gln - Lys - Pro - Val - Ala - Leu - Ile - Asn - Asn - Gln - Phe - Leu - Pro - Tyr - Pro - Tyr -
 40
 Tyr - Ala - Lys - Pro - Ala - Ala - Val - Arg - Ser - Pro - Ala - Gln - Ile - Leu - Gln - Trp - Gln - Val - Leu - Ser -
 50
 Asp - Thr - Val - Pro - Ala - Lys - Ser - Cys - Gln - Ala - Gln - Pro - Thr - Thr - Met - Ala - Arg - His - Pro - His -
 60
 Pro - His - Leu - Ser - Phe - Met - Ala - Ile - Pro - Pro - Lys - Lys - Asn - Gln - Asp - Lys - Thr - Glu - Ile - Pro -
 70
 Thr - Ile - Asn - Thr - Ile - Ala - Ser - Gly - Glu - Pro - Thr - Ser - Thr - Pro - Thr - Ile - Glu - Ala - Val - Glu -
 80
 Ser - Thr - Val - Ala - Thr - Leu - Glu - Ala - Ser - Pro - Glu - Val - Ile - Glu - Ser - Pro - Pro - Glu - Ile - Asn -
 90
 Thr - Val - Gln - Val - Thr - Ser - Thr - Ala - Val-OH
 100
 105 ↓ 106
 110
 130
 136
 140
 148
 150
 160
 169

Figure 4. Primary structure of *Bos* κ -CN B-1P (186). The enclosed amino acid residues are the sites corresponding to the mutational differences in the A variant (186). The arrow indicates the point of attack by chymosin (rennin).

similar reasoning is called β -CN A (f 108-209). As yet, the specific genetic variants of the β -CN fragments (f 1-105), (f 1-107), (f 29-105), and (f 29-107) have not been identified, but their genetic nomenclature is expected to relate in similar manner to the β -CN variant from which they were formed.

κ -Caseins

The κ -casein (κ -CN) family consists of a major carbohydrate-free component and a minimum of six minor components (65, 66, 172, 210, 224, 256, 271, 273). The κ -CN, as isolated from milk, occurs in the form of a mixture of polymers held together by intermolecular disulfide bonds (236). However, Beeby (20) reported free thiol groups after calcium removal by treatment with ethylenediaminetetraacetate (EDTA) or oxalate. He suggested that the native form may be the reduced monomer rather than the disulfide-linked polymers.

The primary structure of the major carbohydrate-free component of κ -CN B-1P (Figure 4) consists of 169 amino acid residues (186): Asp₄, Asn₇, Thr₁₄, Ser₁₂, Ser P₁, Pyroglu₁, Glu₁₂, Gln₁₄, Pro₂₀, Gly₂, Ala₁₅, Cys₂, Val₁₁, Met₂, Ile₁₃, Leu₈, Tyr₉, Phe₄, Lys₉, His₃, Trp₁, and Arg₅ with a calculated molecular

weight of 19, 007. There is still some question concerning the N-terminal pyroglutamyl residue in the native protein because cyclization may occur during isolation (234).


Two genetic variants, designated A and B, are known (189, 223, 270). The κ -CN A-1P differs from the B variant (Table 2) by substitution of a threonine residue for isoleucine at position 136 and an aspartic acid residue for alanine at position 148 (186). The A variant tends to be predominant in most breeds (11). In alkaline gel electrophoresis in the presence of mercaptoethanol and urea, both variants show multiple bands with the A variant possessing the greater mobility (171, 238).

The bond sensitive to chymosin (EC 3.4.23.4) (rennin) hydrolysis occurs between the phenylalanine residue at position 105 and the methionine residue at position 106 (Figure 4) (64, 131, 166, 204). The hydrolytic products are para- κ -casein (residues 1-105) and the macropeptides (residues 106-169). Doi et al. (65) and Vreeman et al. (256) observed para- κ -casein in purified preparations of κ -CN. This is due to a chymosin-like proteolysis subsequent to translation, but more work must be done before concluding that para- κ -casein is a natural constituent of milk rather than a product of storage or of the preparatory process.

Structures of the minor κ -CN components have not been established, and considerable disagreement exists among investigators. Generally, the minor κ -CN are believed to be glycosylated forms of the major κ -CN, but even this has not been confirmed by all investigators. Vreemen et al. (256) concluded from their investigation that, in order of elution from DEAE-cellulose, the adsorbed κ -CN were the major component free of carbohydrate with one phosphate group; a component containing one phosphate and one carbohydrate moiety with a N-acetylneuraminic acid (NeuNAc) residue; a component containing one phosphate group and one carbohydrate moiety with two NeuNAc residues in a branched configuration; a component with two phosphate groups and no NeuNAc; a component with one phosphate and two carbohydrate moieties with two NeuNAc residues each; a component probably containing two phosphate groups and one carbohydrate moiety with two NeuNAc residues; and a component with either two phosphate and two carbohydrates containing two NeuNAc or one

phosphate and three comparable carbohydrate chains. In contrast, Doi et al. (65, 66) concluded from their fractionation of κ -CN on DEAE-cellulose that there are five major and two minor adsorbed components all containing one phosphate group. Of the major components, the first is the major carbohydrate-free component of κ -CN; the second contains 1 mole each of galactose (Gal) and galactosamine (GalNAc) per mole; the third has 1 mole each of Gal, GalNAc, and NeuNAc per mole; the fourth has 3 moles of Gal and 2 moles each of GalNAc and NeuNAc; and the fifth has 4 moles of Gal and 3 each of GalNAc and NeuNAc. The two minor components appear to differ slightly in amino acid content and contain one and two carbohydrate moieties, respectively. Each moiety contains 1 mole each of Gal, GalNAc, and NeuNAc.

Several others have investigated the structure of carbohydrate moieties and their point of attachment to the peptide chain (80, 81, 82, 132, 133, 134, 253, 263, 264). Fournet et al. (81) isolated three oligosaccharides from κ -CN and determined the structures for two:



So far, Thr's at positions 131, 133, and 135 or 136 (81, 82, 130, 132, 134, 142) have been identified as points of attachment of oligosaccharide chains through O-glycosidic linkages. If an oligosaccharide is attached to κ -CN A-1P at Thr 136, the B variant could not contain an oligosaccharide at this position because Ile replaces Thr (Figure 4, Table 2).


Other carbohydrates also have been detected in preparations of κ -CN. Wheelock and Sinkinson (263, 264) found D-glucose (Glu) and D-mannose (Man) in samples of κ -CN from which free sugars were removed. They separated the para- κ -casein from the glycomacropetides after

rennin treatment and found that although the NeuNAc was associated almost entirely with the glycomacropetides, the Man primarily was attached to the para- κ -casein and Gal and GalNAc were in both. More recently para- κ -casein was reported devoid of carbohydrate (81).

The carbohydrate on colostrum κ -CN is more complex and variable than in normal milk (82). Colostrum κ -CN contains almost twice as much carbohydrate as well as an additional amino sugar, N-acetylglucosamine (GlcNAc) (67). Two of the oligosaccharide chains that have been characterized have structures identical to those

(structures 1 and 2) for κ -CN from milk (220, 221). However, four additional oligosaccharides,

which appear to be unique to colostral κ -CN, have been characterized and possess the following structures:



Only Thr's 131, 133, and 135 have been identified as points of attachment for oligosaccharides bound to colostral κ -CN (67).

Because exact structures of minor components of κ -CN are not known, we think that nomenclature of these components cannot be precise at this time. We suggest that they temporarily be identified according to the genetic variant of the major nonglycosylated component and that they be numbered consecutively according to their increasing relative electrophoretic mobility in alkaline gels in the presence of mercaptoethanol (273) starting as κ -CN A-1 or κ -CN B-1. More conclusive results are needed before nomenclature of the κ -CN's can be finalized.

WHEY PROTEINS

The term "whey proteins" has been used to describe the group of milk proteins that remain soluble in "milk serum" or whey after precipitation of caseins at pH 4.6 and 20°C. Traditionally, β -lactoglobulin, α -lactalbumin, serum albumin, immunoglobulins, and proteose peptone fractions have been considered to be the major characterized components of this fraction. However, recent findings that proteose peptone components 5, 8-slow, and 8-fast are actually fragments of β -CN (7, 8, 74) and that

other heat stable polypeptides are present (197) raise questions concerning the utility of this term. Based on current knowledge, the term "whey protein" should be used only in a general sense to describe milk proteins soluble at pH 4.6 and 20°C. Commercial products termed whey protein isolates or concentrates are obtained from cheese manufacture at higher pH's and will contain intact caseins as well as their proteolytic products such as macropeptides and proteose peptone fraction. Individual families, such as β -lactoglobulin, α -lactalbumin, and serum albumin, should be classified according to homology with the primary sequence of their amino acid chains. Polyacrylamide or starch gel electrophoresis still can be used to characterize and identify individual members of each family (238). The characterization of immunoglobulins, proteins not unique to milk, continues to be based upon immunochemical as well as physical and chemical similarities with human and other mammalian immunoglobulins.

β -Lactoglobulin

β -Lactoglobulin B (β -LG) consists of 162 amino acid residues with the following composition: Asp₁₀, Asn₅, Thr₈, Ser₇, Glu₁₆,

Gln₉, Pro₈, Gly₄, Ala₁₅, Cys₅, Val₉, Met₄, Ile₁₀, Leu₂₂, Tyr₄, Phe₄, Lys₁₅, His₂, Trp₂, and Arg₃, and a calculated molecular weight of 18,277. The primary structure (Figure 5) has been determined by Braunitzer and coworkers (35, 36, 85, 86). Several corrections to the original sequence have been made and are included in the primary structure shown for β -LG (Figure 5). Residues 155 and 156 have been changed from leucine-glycine to glycine-leucine (100, 207), and positions 84 and 87 are occupied by isoleucine and leucine instead of leucine and isoleucine (207). However, the proposal that residue 11 is asparagine instead of aspartic acid (100) has not been substantiated (25, 207) and still is shown to be aspartic acid (Figure 5) until additional information becomes available. The one free thiol group that occurs in each molecule of β -Lg appears to be distributed equally between positions 119 and 121 (36, 180, 182). When the thiol occurs at residue 119, a disulfide bridge occurs between residues 106 and 121, and between residues 106 and 119 when the free thiol is at position 121. In either situation, the other disulfide bridge always occurs between positions 66 and 160 (Figure 5).

Seven genetic variants of β -LG are known (12, 24, 25, 40, 100, 199). Positions of amino

acid substitutions have been determined for five of these variants, A, B, C, D, and E (Figure 5, Table 2). Yak (*Bos grunniens*) milk contains only a single type of β -LG with an electrophoretic mobility similar to bovine β -LG D (100). However, this β -LG differs from the bovine B variant (Figure 5) by substitution of glycine for glutamic acid at 158 (100) instead of glutamine for glutamic acid at 45 as in the D variant (Table 2). Grosclaude et al. (100) suggested that this variant be labeled β -LG Dyak. However, because we wish to avoid subscripts and superscripts, we suggest that this variant in yak milk be called, in order of its discovery, β -LG E (Table 2). Three additional genetic variants of β -LG (E, F, and G) have been observed in the milk of Bali (Banteng) cattle, *Bos (Bibos) javanicus* (25). Through a combination of amino acid analysis, tryptic peptide maps, and amino acid analyses of different peptides, Bell et al. (25) showed that β -LG E in Bali milk appears to possess the same glycine/glutamic acid substitution at position 158 as the β -LG Dyak we now suggest be called β -LG E. The β -LG F appears to differ from the bovine B variant by this same substitution at 158 as well as substitution of serine for proline at 50 and tyrosine for aspartic acid at either 129 or 130. The β -LG G also appears to differ from




Figure 5. Primary structure of *Bos* β -LG B (35, 36, 85, 86). The locations of the SH-group and the disulfides are deduced by applying the observations of McKenzie et al (180, 182) to the most recent primary sequence of Braunitzer et al (85, 86). The SH-group is assumed to exist in a 50:50 distribution between positions 119 and 121 with the -SS-bridge location depending upon the position of the SH-group. The enclosed amino acid residues are the sites corresponding to the differences in the genetic variants.

the B variant by substitution of glycine for glutamic acid at position 158 as well as methionine for isoleucine at position 78 (Table 2). However, the exact locations of amino acid substitutions in β -LG F and G still need to be verified by sequencing of the amino acid chains.

A report of this Committee (267) mentioned an additional variant of β -LG occurring rarely in milk of Droughtmaster cattle (24). This variant originally was reported to be identical to the A variant except for the presence of covalently-attached carbohydrate. Carbohydrates present are NeuNAc, GlcNAc, GalNAc, Man, and Gal in the proportions 1.0:3.4:1.9:1.9:8 (24). This glycosylated form should not be considered a genetic variant of β -LG, because it does not appear to occur through point mutation or deletion but rather through posttranslational modification. Recently this variant was reported to differ from β -LG A by substitution of asparagine for aspartic acid at residue 28 (25), the point of attachment for the carbohydrate moiety. If this substitution can be established, then this form of β -LG in Droughtmaster milk is a true genetic variant. In the meantime, we suggest that the term β -LG Dr be retained temporarily until the nature of the variation is established.

α -Lactalbumin

The interaction of α -lactalbumin (α -LA) with galactosyltransferase to produce lactose has been the subject of several reviews (39, 70,

117, 136, 137). The complete amino acid sequence of bovine α -LA has been determined (Figure 6) by Brew et al. (37, 38, 255). As shown, the B variant consists of 123 amino acid residues with the composition: Asp₉, Asn₁₂, Thr₇, Ser₇, Glu₈, Gln₅, Pro₂, Gly₆, Ala₃, Cys₈, Val₆, Met₁, Ile₈, Leu₁₃, Tyr₄, Phe₄, Lys₁₂, His₃, Trp₄, and Arg₁, and a calculated molecular weight of 14,175. All of the half-cysteine residues are connected by intramolecular disulfide bonds (Figure 6). Studies involving metal analyses and effects of EDTA on denaturation indicate that α -LA binds calcium (118).

Three genetic variants, A, B, and C, are known. Only the B variant has been observed in milk from Western cattle, whereas both A and B occur in milk from African Fulani and African and Indian Zebu cattle (10, 29). The α -LA A differs from B in the substitution of a glutamine residue for arginine at position 10 (Table 2) (93). The α -LA prepared from milk of Bali cattle migrates more slowly in alkaline gel electrophoresis than bovine α -LA B (23). Although designated as α -LA C (23), the amino acid substitutions that differentiate this variant have not been established.

Minor forms of bovine α -LA have been reported and some of these contain carbohydrates (13, 16, 17, 23, 93, 124). Aschaffenburg and Drewry (13) observed a faster moving component in an α -LA preparation on paper electrophoresis at pH 8.6. This protein had




Figure 6. Primary sequence of *Bos* α -LA B (37, 38, 255). The disulfide bridges in the molecule are as indicated between positions 6 and 120; 28 and 111; 61 and 77; and 73 and 91. The enclosed amino acid residue is the site corresponding to the difference in genetic variant A.

the same amino acid composition as α -LA but contained one additional hexosamine residue per molecule (93, 94). Barman (16) prepared a glycosylated form of α -LA containing Man, Gal, fucose, GlcNAc, GalNAc, and NeuNAc in proportions 4.1:1.4:1.0:3.1:1.1:64. Hopper and McKenzie (124) examined minor components of α -LA A and B and found one component with faster electrophoretic mobility than α -LA in starch gels at pH 7.7 and two with slower electrophoretic mobilities. The fast component appears to have one less amide group than α -LA. The faster of the two slow components contains NeuNAc, but the slower one does not. Although α -LA possesses four disulfide bonds (255), Barman (17) reported isolation of a minor fraction (5%) containing only three disulfide bonds. All of these minor components still retain their lactose synthetase activity. More information, therefore, is needed before nomenclature of these minor components of α -LA can be established. In the interim, use of Arabic numbers to differentiate by electrophoretic mobility is suggested.

Serum Albumin

Serum albumin (SA) prepared from milk is physically (203) and immunologically (62) identical to blood SA, the major protein of plasma; the protein prepared from blood plasma has been studied extensively. Since the last report of this committee (267), a complete amino acid sequence of bovine SA has been determined (Figure 7) by Brown (43, 44). The original sequence presented by Brown (43) contained a gap of three (43) or four (44) residues extending from 400 to 403; these residues were identified as Gly-Phe-Gln-Asn by Reed and coworkers (214) and are included in the sequence in Figure 7. The sequence presented by Brown (44) assumed lysine at residue 400 as in human SA (22, 184). However, Reed and coworkers (214) found this residue in *Bos* was glycine rather than lysine. *Bos* SA consists of 582 residues with the composition: Asp₃₉, Asn₁₂, Asx₃, Thr₃₄, Ser₂₈, Glu₅₉, Gln₁₉, Glx₁, Pro₂₈, Gly₁₆, Ala₄₆, Cys₃₅, Val₃₆, Met₄, Ile₁₄, Leu₆₁, Tyr₁₉, Phe₂₇, Lys₅₉, His₁₇, Trp₂, and Arg₂₃ and a calculated molecular weight of 66,267. Serum albumin has 17 intramolecular disulfide bonds and only one free sulfhydryl group, which occurs at residue 34 (Figure 7). The sequence and disulfide bridges shown

do not indicate the structural features of loops and domains proposed by Brown (44). The molecule is visualized as having three major domains, each consisting of two large double loops and a small double loop and appears to be approximately in the shape of 3:1 ellipsoid. The N-terminal region is believed to be more compact than the C-terminal region (198). The domains are dissimilar in hydrophobicity, net charge, and ligand binding sites (198). The appearance of several bands in isoelectric focusing patterns indicate that considerable microheterogeneity may exist (232).

Immunoglobulins

Like SA, immunoglobulins are not unique to lacteal secretions. Except for secretory component (SC) and the complex called secretory immunoglobulin A (SIgA), which it forms with immunoglobulin A (IgA), all immunoglobulins in milk are normally in serum, although concentrations of each in colostrum and milk may vary (34, 52, 53, 168, 205). However, because of their extreme heterogeneity, immunoglobulins differ significantly from other serum proteins. Thus, physical and chemical parameters routinely used to characterize and identify most other milk proteins are only of limited value for immunoglobulins. Their identification must be based primarily upon immunochemical criteria and crossreactivity with reference immunoglobulins that are primarily of human origin. A system of nomenclature already has been proposed for the bovine immunoglobulins (1, 2, 60) based upon a system originally proposed for those of human origin by the World Health Organization (61) and revised (191). Our recommendations for nomenclature of the immunoglobulins in bovine milk, as well as their structural components and closely related proteins, continue to be based upon these systems (1, 2, 60, 61, 191).

Of the five classes of immunoglobulins in mammals, four (IgG, IgA, IgM, and IgE) have been identified in bovine milk. All classes of immunoglobulins exist as either polymers or monomers of a basic unit composed of four polypeptide chains. In all species in which immunoglobulins have been studied extensively, the monomeric form consists of two identical heavy (molecular weight from 50,000 to 70,000, depending upon class) and two identical light (molecular weight about 20,000) poly-




Figure 7. Primary sequence of *Bos* SA (43, 44). The 17 disulfide bridges are indicated by solid lines; the only free sulfhydryl group occurs as residue 34.

peptide chains that are linked covalently by disulfide bonds. All evidence indicates that basic structures for the various classes of immunoglobulins in *Bos* are similar to those of human origin (49, 50, 55, 148, 155). The N-terminal half (first 107 to 115 amino acid residues) of each pair of identical light polypeptide chains is called the variable region because of its

highly variable amino acid sequence. Because the sequence of the C-terminal halves are more similar, they are called the constant region. Each pair of identical heavy polypeptide chains also contains a variable (first 107 to 115 residues depending on the variable region subgroup) and a constant (310 to 500 amino acid residues at C-terminus) region. Variable regions of both heavy and light polypeptide chains are responsible for antigen binding, whereas functions of complement fixation, membrane transport, catabolism, and mediation of immediate-type hypersensitivity all have been attributed to the constant region of the heavy polypeptide chains (90). The lower portion of this region generated by proteolysis (51) has been isolated and crystallized and is termed the crystalline fraction (Fc).

Each class of immunoglobulin differs by the sequence of amino acids in the constant region of their heavy polypeptide chains. These differences can be recognized primarily by use of specific antisera to these class-specific determinants. Greek letters, corresponding to Latin letters used to designate each class of immunoglobulins, are assigned to each pair of heavy polypeptide chains that account for these differences (e.g., γ for heavy chains of IgG) (49, 90). When differences in amino acid sequences and, consequently, antigenic determinants, are more subtle, the difference is one of subclass. Numerical subscripts are used with Greek letters to designate these heavy polypeptide chains (e.g., γ_1 for heavy chains in IgG₁). The term allotype is used to designate genetic variations within species similar to blood group antigens.

Light polypeptide chains also can be differentiated by variations within their constant regions. Two such types of light chains, called κ and λ , occur in all classes of immunoglobulins (90). Although both types of light chains also occur in bovine immunoglobulins, the proportion of λ type light chains is higher (122).

Immunoglobulin G (IgG) is the principal class in bovine colostrum and milk. The IgG exists primarily as a monomer of the basic four polypeptide chain unit consisting of two identical heavy (γ) and two identical light (λ or κ) polypeptide chains. The two major subclasses in milk and colostrum of *Bos* are IgG₁ and IgG₂. These subclasses can be distinguished antigenically by determinants in the Fc portion

of the heavy (γ_1 and γ_2) polypeptide chains (78). Common antigenic determinants are also in these same constant regions (209; K. Nielson, personal communication). Even though bovine serum contains nearly equivalent amounts of IgG₁ and IgG₂, colostrum and precolostral secretions typically contain 15 to 20 times as much IgG₁ as IgG₂ (113). After transition from colostrum to milk, IgG₁ levels still remain 4 to 7 times greater than IgG₂ (113). Elevated IgG₂ has been reported in milk during inflammation of the gland, the source of which has been discussed elsewhere (57, 170, 257, 274). Low molecular weight forms of IgG have been found in bovine milk and colostrum (21, 51, 92). Although some have been similar to the F(ab)₂, Fab, and Fc fragments, which are generated by proteolysis of IgG with pepsin and papain (51), additional information is needed before their nomenclature can be recommended.

The considerable charge heterogeneity between IgG₁ and IgG₂ provides an excellent example for the necessity of using immunological criteria for identification and characterization of immunoglobulins. Ion-exchange chromatography can be used routinely to obtain relatively pure preparations of IgG₁ and IgG₂. Usually, IgG₂ does not bind or is only weakly bound by DEAE- or QAE-cellulose, whereas IgG₁ is removed through use of salt. However, in some cattle the extreme charge heterogeneity IgG₁ and IgG₂ results in overlap of the chromatographic peaks to such an extent that purification, characterization, or both, based solely upon ion-exchange behavior is virtually impossible (54, 55). Only immunological crossreactivity with authentic bovine IgG₁ and IgG₂ can be used for identification.

The IgA in bovine lacteal secretions exists primarily as a dimer of the basic four polypeptide chain unit (58, 68, 96), although some higher aggregates and some apparent degradation products can be found both in milk and colostrum. The basic polypeptide chain unit of IgA consists of two identical heavy (α) and two identical light (λ or κ) chains. Bovine IgA concentrations are elevated in colostrum with concentrations dropping rapidly throughout lactation. Total output of IgA by the mammary gland appears to remain nearly constant (2.5 g/day) throughout lactation; changes in volume greatly influence actual

concentration during various stages of lactation (113).

In addition to existing as a dimer in lacteal secretions, IgA also occurs in a complex called SIgA (59, 152, 167, 212, 247), which consists of 1 mole of dimeric IgA covalently attached to 1 mole each of SC and J-chain. Bovine SC, which was isolated first in free form from milk and called glycoprotein a (50, 56, 108), has a molecular weight of 79,000 calculated from its amino acid content (160). Both IgA and SC have been isolated and characterized (50, 68, 160, 168, 169, 206, 254) and are homologous to human IgA and SC (55, 160, 167). Heterogeneity in the SC fraction may be due in part to varying degrees of glycosylation (160) and in part to occurrence of proteolytic fragments known to form during storage of SC (208). More information concerning the nature of this heterogeneity is needed before nomenclature can be recommended. Bovine J-chain has a molecular weight of 16,500 and contains about 10 thiols/mole.

Immunoglobulin M (IgM), which has a sedimentation constant of 19 s (96, 158), usually exists as a pentamer of the basic four polypeptide chain unit. Each mole of pentameric IgM in milk appears also to contain 1 mole of covalently-bound J-chain as well as about 1 mole of noncovalently attached J-chain (155). Each basic four polypeptide chain unit of IgM consists of two identical heavy (μ) and two identical light (λ or κ) polypeptide chains. Bovine IgM, which has been purified from milk by a variety of methods (79, 251), is antigenically homologous to human IgM (183) as well as similar to its counterparts in other species. The IgM concentration in milk averages

about .1 mg/ml (113, 251) and is elevated in colostrum (about 6 mg/ml) with gradual decline throughout lactation. Output of IgM in mid-lactation milk is about 1 g/day, about half that produced during early lactation (113).

A protein that possesses reagenic activity and does not belong to IgG, IgA, or IgM classes is transmitted to suckled calves through colostrum (26, 115). This protein generally has been accepted to be IgE and crossreacts with human IgE (192). The molecular size of this protein falls between IgG and IgM (115, 261), which is consistent with the behavior of IgE in other species. Additional information is needed to characterize more fully this bovine immunoglobulin.

β_2 -Microglobulin

β_2 -Microglobulin (β_2 -m) first was prepared from bovine milk by Groves and coworkers (105, 107) as lactollin. This minor milk protein is mentioned in this report since Groves and Greenberg (111, 112) showed that lactollin is bovine β_2 -m. Human β_2 -m is part of the histocompatibility complex in man and is thought to be necessary for native conformation and antigenic activity of the latter (162). Human β_2 -m shows homology to the constant regions of human IgG heavy and light chains, especially CH3 (200).

Bovine β_2 -m concentrations were higher in colostrum than in milk (105, 107). Measurements by competitive immunoassay revealed concentrations as high as 6.0 μ g/ml in colostrum and about 4.0 μ g/ml in mature milk (208). The amino acid sequence of bovine β_2 -m has been completed (112) and is in Figure 8. Bovine




Figure 8. Primary sequence of *Bos* β_2 -m determined by Groves and Greenberg (112). β_2 -m has been shown to be identical with a minor milk protein previously identified as lactollin (111, 112).

β_2 -m consists of 98 amino acid residues with the composition: Asp₇, Asn₄, Thr₂, Ser₉, Glu₇, Gln₅, Pro₉, Gly₃, Ala₁, Cys₂, Val₅, Ile₆, Leu₈, Tyr₆, Phe₄, Lys₉, His₄, Trp₂, and Arg₅ with a calculated molecular weight of 11,636. Several physical properties of β_2 -m have been determined recently (159).

PROTEIN OF MILK FAT GLOBULE MEMBRANE

According to King (150), first mention that fat droplets dispersed in milk are surrounded by a thin membrane was by Ascherson in 1840. For over 120 yr considerable controversy existed over both the origin and nature of this membrane (45, 150). The first report published by this Committee in 1956 (129) defined proteins associated with this membrane as:

a complex of proteins and enzymes including lipoproteins, alkaline phosphatase, and xanthine oxidase adsorbed on the surface of milk fat globules. This material (or at least a fraction thereof) can be isolated by washing cream and churning.

The nomenclature of milk fat globule membrane proteins was not mentioned in subsequent reports of this Committee (48, 218, 246, 267), primarily because these polypeptides had not been characterized sufficiently. However, during the past 10 yr, a more definitive picture of the milk fat globule membrane (MFGM) and its associated polypeptides has evolved through increased knowledge of biological membranes (230) and of the mechanism of synthesis and secretion of milk (4, 146, 147, 196, 269). Fat droplets form intracellularly and become enveloped progressively in apical plasma membrane as they are secreted into the alveolar lumen. Thus, proteins associated with the MFGM are derived in part from the apical plasma membrane and in part from the cytosol through adherence to fat droplets or to the internal side of the plasma membrane (89, 269). Thus, a concise catalogue of all proteins associated with MFGM may not be possible.

In keeping with history, we propose that an operational definition of MFGM be developed until associated proteins can be characterized with respect to biological function or primary sequence of their amino acid chain. This definition must be broad enough to allow for inclusion of new discoveries and observations

but, as for skim milk proteins, will not include enumeration of enzyme activities associated with MFGM (4, 45, 196).

Singer (230) proposed use of the terms peripheral or integral to describe the degree of association of proteins with various biological membranes such as MFGM. An integral protein is associated closely with the membrane; that is, it actually may be imbedded in or transverse the membrane and is not washed off easily by a variety of reagents. In contrast, a peripheral protein is associated loosely with the membrane and may be removed easily by washing. The association of α -LA with galactosyltransferase in the Golgi apparatus of lactating mammary gland provides an excellent example. The α -LA is considered a peripheral membrane protein because of its association with galactosyltransferase, whereas the enzyme is an integral component of the Golgi apparatus membrane (230). Although we are dealing with a true biological membrane, our consideration of its proteins is within the context of milk proteins. This rules out peripheral proteins as components of MFGM. Rather, the definition of integral protein appears to suit our needs better at this time. Moreover, most researchers working with MFGM have employed some type of wash to remove skim milk proteins from fat globules (6, 116, 154, 239) and, thus, operationally have excluded α -LA and other peripheral skim milk proteins.

In addition to recommending a method for washing lipid droplets, an operational definition of MFGM proteins also should include history of the milk, method of churning to liberate the membranes, method of harvesting the membranes, and method of separation and characterization of the protein components. Given this operational definition, a provisional nomenclature can be devised to provide better communication among researchers concerned with MFGM proteins.

History of Milk

Because cooling of milk prior to preparation of MFGM leads to changes in morphology (269), crystallization of triglycerides (45), and loss of phospholipids, the use of warm milk has been recommended widely. However, Kobylka and Carraway (154) showed that prior chilling of the milk actually has little effect on overall distribution of major membrane proteins.

In the specific case of MFGM proteins, however, overriding consideration must be given to the presence of proteolytic enzymes such as plasmin. Plasminogen is associated with MFGM preparations (121), and fragments of membrane proteins have been found elsewhere in milk, particularly in the proteose peptone fraction (145). Preparation of MFGM from milks that have aged overnight (5) may lead to generation of artifacts. Thus, the use of fresh, warm, raw milk for the study of MFGM is recommended strongly. However, almost all market milks are subjected to cold storage, and the functional properties, which make dried buttermilk a useful commodity, may be the result of proteolytic action.

Washing of Cream

Although many techniques have been employed to harvest fat globules from milk, centrifugation followed by washing is the most widely used. Fresh warm milk is skimmed conveniently by centrifugation at 5,000 to 10,000 $\times g$ for 10 to 20 min at room temperature or higher (153, 154, 178). Large scale preparations generally employ a laboratory separator operated at 40°C (6, 116, 144, 196). In general, cream then is standardized to approximately 30% fat and washed. Deionized water (18, 116, 144), buffered saline (89, 193), and buffered sucrose with (178, 231) or without added Mg^{2+} (6) have been used to wash lipid droplets. Although use of sucrose $\pm Mg^{2+}$ appears to offer some advantage in stabilizing enzymatic activities (4, 45, 239), electrophoretic patterns of MFGM prepared with the various washing media appear to be qualitatively similar (Figure 9). Variations in washing media appear primarily to affect distributions of major membrane proteins. A similar conclusion can be reached by an examination of the literature. Finally, three washes of the fat globules appear to be optimal (179, 239).

Churning

Rupture of fat globules by physical or mechanical means is necessary to liberate membranes. This has been accomplished effectively by freeze thawing (153, 154, 231), blending (6, 18, 178), or by use of a rotating shaker (173, 239). The question of cooling is raised again. It is difficult to churn warm cream

effectively. Cooled cream requires less mechanical energy to churn; however, some proteins that are associated loosely with the membrane through hydrophobic interactions could be lost.

Recovery of Membranes

The best method for separating MFGM from buttermilk and fat is by warming the churned material to 37 to 40°C followed by centrifugation of the warmed suspension (6, 154, 178). Crude membrane preparations are pelleted during centrifugation. Times and centrifugal forces vary; however, it appears that standardization at 100,000 $\times g$ for 1 h should be recommended. Washing and repelleting can be employed. In large scale operations, butter oil is




Figure 9. The SDS gel electrophoresis of proteins from milk fat globule membrane (MFGM) prepared by methods of 1) Carraway et al. (153, 231), 2) Anderson et al. (6), 3) Mather and Keenan (178), 4) Nielson and Bjerrum (193), 5) Brunner et al. (45), and 6) Jarasch et al. (125). The gel is a 4 mm slab with 4% and 15% acrylamide in the stacking and separating gels, respectively. The method is essentially that of Laemmli (161). Lane S contains α -LA, trypsin inhibitor, carbonic anhydrase, ovalbumin, SA, and phosphorylase b (bottom to top), which were used as molecular weight markers.

removed by use of a laboratory separator at 40°C, and the membrane may be floated to the top of the buttermilk with (NH₄)₂SO₄ (18, 116, 144).

Electrophoretic Separations

The most universally accepted method for characterization of membrane proteins employs electrophoresis in the presence of the detergent sodium dodecyl sulfate (SDS). Thus, the extraction procedure used prior to SDS gel electrophoresis is an integral part of any operational definition for membrane proteins. Although several procedures have been developed for extracting membrane proteins (190, 259, 272), the most successful, and apparently artifact-free, method (201) involves incubation at 37°C for 30 min at alkaline pH with 10% (wt/vol) 2-mercaptoethanol. (Less reductant (272) and shorter incubation times work well with soluble proteins.) Additionally, the extraction medium must contain 1.4 g SDS/g protein, and the concentration of SDS must be at least 1% (wt/vol) (272).

Most workers (18, 151, 173, 178, 229) have used SDS gel electrophoretic procedures to characterize MFGM proteins. These are modifications of the Weber and Osborn (259) continuous buffer gels and give reasonably good resolution of membrane proteins. Anderson et al. (5) employed acrylamide gradient gels, but this method has not been used as widely. Thus far, in all the SDS gel electrophoretic systems attempted, fine division of MFGM proteins has been obtained only in the most carefully done work (178, 229). However, use of a discontinuous gel electrophoretic system, such as that described by Laemmli (161), routinely produces excellent resolution of the fine structure of complex mixtures. This method appears to hold great promise for successful resolution of MFGM proteins as well (19, 177). However, discontinuous gel electrophoresis in SDS must be used with precautions described by Wyckoff et al. (272). These include modifying the Laemmli procedure (161) to eliminate SDS from the gel buffers, reducing SDS in the upper buffer to .03% (wt/vol), and reducing SDS in

the total sample load to 150 µg. With these modifications, good resolution of MFGM proteins can be obtained (Figure 9) by the stacking and 10 to 15% acrylamide separating systems of Laemmli (161). This system currently is packaged commercially as Sage by Miles, Inc.²

Definition of Milk Fat Globule Membrane Protein

The Committee recommends the following criteria to serve as an operational definition of MFGM proteins. First, membrane proteins must be associated with the cream fraction obtained from raw uncooled whole milk and remain with the cream after three washings of the fat globules in an aqueous medium. Second, after churning of cream by mechanical means and removal of butter oil at 37 to 40°C, membrane proteins must remain associated with the buttermilk fraction. Third, the protein must coisolate with the membrane fraction prepared from buttermilk either by centrifugation for at least 1 h at 100,000 × g at 4°C or by salting out with 2.2 M (NH₄)₂SO₄. Finally, the protein should be extractable with 1% SDS (1.4 g SDS/g protein) and 10% 2-mercaptoethanol and be visualized by staining with either Coomassie blue, periodate acid Schiff reagent or both after SDS gel electrophoresis.

Suggested Nomenclature for Milk Fat Globule Membrane Proteins

When MFGM preparations and molecular weight standards containing α-LA, ovalbumin, and phosphorylase b are examined by the modified Laemmli (161) method described for SDS gel electrophoresis, membrane proteins can be separated into four distinct zones based on their electrophoretic mobilities (Figure 9). Zone A lies between the stacking gel-separating gel interface and phosphorylase b, Zone B between phosphorylase b and ovalbumin, Zone C between ovalbumin and α-LA, and Zone D represents all lower molecular weight proteins that may be true membrane proteins or may be proteolytic products that are not resolved readily and migrate at the "front" of the 15% gels. The majority of proteins associated with MFGM fall into Zones A and B described previously. Because higher percent acrylamide gels require special care, the Committee

²Reference to brand or firm does not constitute endorsement by this committee over others of a similar nature not mentioned.

recommends that the standard gel contain 15% acrylamide. However, because elucidation of Zone A proteins may require lowered content of acrylamide in gels, the nomenclature will allow designation of the percent acrylamide in gels used to characterize the membrane proteins. Changes in percent acrylamide of gels may allow bands to cross over because glycoproteins, in particular, provide anomalous results (5). This situation may be dealt with as the need arises.

The second notation to be used in describing a MFGM protein is the "apparent" molecular weight as interpolated by comparison with standard gels containing molecular weight markers in addition to α -LA, ovalbumin, and phosphorylase b. Finally, the third portion of the nomenclature describes whether the protein stains with Coomassie blue (C), periodate acid Schiff (S), or both after separation by gel electrophoresis. The complete description of a hypothetical protein is:

MFGM-A₁₅-127, C, S

where MFGM indicates that the protein satisfied the operational definition for a MFGM protein, A₁₅ designates the zone in a 15% Laemmli (161) gel, 127 is the "apparent" molecular weight in k-daltons, and C, S designates that the protein is both Coomassie blue and periodic acid Schiff positive. We recognize the prerogative of the investigators who first describe a protein in detail to provide a trivial name, but we suggest that the nomenclature described be observed as well. One such trivial name that recently has been suggested is butryophilin (87).

NOMENCLATURE OF MILK PROTEINS OF SPECIES OTHER THAN THOSE OF THE GENUS *BOS*

Gel electrophoresis, both in the presence and absence of SDS, has been used to characterize proteins in milks of several species that do not belong to the genus *Bos*. In most cases, the gel electrophoretic patterns are reasonably complex and indicate that milks of these species contain numerous distinct polypeptides. Furthermore, gel electrophoretic techniques often are used to monitor isolation of these milk proteins and to determine purity of a particular preparation. In either situation, a temptation exists to assign names to bands in the electrophoretic patterns obtained from these milks, or to polypeptides

actually purified from them, based upon similarities of their electrophoretic mobilities to proteins in bovine milk. However, electrophoretic methods are not good indicators of interspecies protein homology. Classification of a milk protein as a homolog of one of those in *Bos* on the basis of electrophoretic similarity may produce errors. Instead, in this section, we recommend a system of nomenclature to be used for naming proteins obtained from milks of species not belonging to the genus *Bos*. Although this represents a major deviation from the scope of previous reports, we think it is justified by the past tendency of investigators to base nomenclature of milk proteins of other species on that already established for bovine milk.

Nomenclature of Species

We recommend that at first mention of a milk protein in a publication, the mammalian species from which it was obtained be designated by its Latin binomial and vernacular names. Thereafter in the publication either Latin or vernacular nomenclature for the species may be used. A check list of Latin binomial and English vernacular names for mammals is available (123).

Nomenclature of Polypeptide Chain Families

Because species and subspecies within the genus *Bos* can interbreed freely, each of their families of milk proteins can be considered a common pool of genetic polymorphs just as though they were intraspecies variants. Thus far, some milk proteins of Indian cattle (*Bos indicus*), yak (*Bos grunniens*), and Bali cattle (*Bos banteng*) have been isolated and characterized and are included in the tabulation of bovine variants (Table 1). Proteins that may be isolated from milks of other species of the genus *Bos* and characterized should be treated in similar fashion.

We recommend that milk proteins obtained from species not belonging to the genus *Bos* be named, as far as possible, as homologs of the families of proteins known to occur in milk of *Bos*. This recommendation is consistent with the current tendency to name milk proteins from these other species after their bovine counterparts. Names of the principal families of proteins in the milk of *Bos* have been given along with suggested abbreviations (Table 1). Each name,

some containing Greek letter prefixes, designates a family of homologous nascent polypeptide chains. Milks of other species contain recognizable homologs of many, but not necessarily all, proteins in bovine milk. For example, milks of humans, guinea pigs, rats, and mice do not contain β -LG (126). However, milks of some other species contain significant amounts of proteins such as lactoferrin, transferrin, and lysozyme, which occur only in extremely low concentrations in bovine milk. Additionally, rat and mouse milks contain, respectively, a group of phosphoproteins, designated P proteins (181), and a lipoprotein, called whey acidic protein (202), for which corresponding homologs have not been identified in bovine milk. When this situation occurs, assignment of nomenclature is left to the discretion of the investigator who is responsible for establishing the uniqueness of that particular milk protein.

Some may question whether milk proteins obtained from either of the two extant species of bison (*Bison bison* and *Bison bonasus*) should be treated as genetic variants or species homologs of bovine milk proteins. Although species belonging to *Bos* and *Bison* interbreed, not all crossbred progeny are fertile. Thus, for the present, we recommend that bison milk proteins be considered homologs, rather than genetic variants, of proteins in bovine milk. To our knowledge, no bison milk proteins have been isolated and characterized. Certainly proteins in the milks of Asiatic and African buffalo (*Bubalus* and *Synceros*) should not be considered genetic variants of *Bos* proteins as members of these two genera do not interbreed with any species belonging to *Bos*.

Nomenclature Assignment

The major difficulty in assigning nomenclature to proteins obtained from milks of other species is in determining their homology with bovine milk proteins. In biological terminology, homology implies divergence from a common ancestor. Although homology can be inferred from similarity in biological function, this characteristic alone should not be considered a safe indicator, as it could have arisen by convergence from different ancestral lines. Although homology also can be inferred through immunological crossreactivity, the most definitive criterion is congruence of amino acid sequence (268).

When a new protein first is isolated from milk of another species, we recommend that it be assigned a provisional name until its homology with a bovine milk protein can be established. If the protein is clearly a casein or whey protein, it can be designated so in the provisional nomenclature. If the protein does not fall clearly into either of these categories, it simply can be called "milk protein". Roman numerals (which are not used anywhere in the permanent nomenclature described in this paper) can be used to designate any fractions that result from application of electrophoresis, gel filtration, or ion-exchange chromatography. Examples of this provisional nomenclature would be domestic cat (*Felis catus*) casein II, raccoon (*Procyon lotor*) whey protein IV, and golden hamster (*Mesocricetus auratus*) milk protein V.

Because the amino acid composition of a protein is usually available long before the primary sequence of the amino acid chain, the ability to determine the existence of homology between two proteins through comparison of their amino acid compositions would be advantageous. The $S\Delta Q$ characteristic suggested by Marchalonis and Weltman (176) is suitable for this purpose:

$$S\Delta Q = \sum_j (X_{ij} - X_{kj})^2$$

where i and k identify proteins being compared and X_j is the content in moles/100 moles of the amino acid j . If two proteins have $S\Delta Q$ less than 50, we recommend that they be considered homologous until that time at which their primary sequences can be determined and compared. Homologous protein pairs generally have $S\Delta Q$ well below 100 (176). If the protein pairs share common biological function as well as $S\Delta Q$ below 50, homology can be assigned with more certainty. However, similarity in primary sequence of the polypeptide chains still must be retained as the final test for homology between two proteins. In general, two proteins can be considered homologous if their total primary sequences do not differ by more than 50% (63) or if their N-terminal sequences of 20 amino acids do not differ by more than 5 substitutions.

We recommend that genetic polymorphs of each homologous protein within species be designated by placing capital Latin letters after the family name as is currently the practice for bovine milk proteins. Latin letters should be used only to designate genetic variants that have been shown to differ in the substitution of one or more amino acids in the nascent polypeptide chain. These letters should not be used to denote differences in electrophoretic mobilities that might just as easily be due to differences in posttranslational modification. Furthermore, situations such as have occurred with bovine β -CN's A¹, A², and A³ should be avoided. As described, these three genetic variants migrate as a single band in alkaline gels but are resolved as three bands in acid gels. We recommend that no subscripts or superscripts be used. Latin designators should be assigned in chronological order of discovery and characterization of the genetic variant. If only a single genetic variant of a protein is found initially, it immediately should be assigned the designator "A" to preclude possible confusion in the literature when additional variants are discovered.

Posttranslational Modification

We recommend placement of an Arabic number after the Latin designator to differentiate between polypeptide chains that differ as a result of varying degrees of posttranslational phosphorylation, glycosylation, or deamination. As suggested for the various glycosylated and phosphorylated forms of bovine κ -CN, proteins should be numbered consecutively in order of electrophoretic mobility in alkaline gels. When the degree of phosphorylation is known, the Arabic number, along with the letter P, will be used to indicate the actual number of phosphate moieties attached to each polypeptide chain. Thus, two goat caseins that contain 6 and 5 phosphates per molecule and previously were called β_1 - and β_2 -caseins (217) now would be designated goat (*Capra hircus*) β -CN A-6P and goat (*Capra hircus*) β -CN A-5P, respectively.

We also recommend that proteins present as a result of proteolytic cleavage in milks of other species be named as fragments of the parent polypeptide from which they are derived. The fragment will be designated by enclosing the

sequence position number of the first and last amino acid residue of the fragment in parentheses preceded by the letter "f". Thus far, only bovine milk definitely has been shown to contain proteins that are proteolytic derivatives. For example, as discussed, γ_1 -casein A² is in bovine milk as a result of hydrolysis of β -CN A² at Lys 28 by plasmin. Thus, the nomenclature of γ_1 -casein A² has been changed to β -CN A²-1P (f 29-209). Proteolytic fragments in milks of other species should be named in similar fashion when sufficient information is available.

Characteristics of proteins isolated from milks of species other than *Bos* have been summarized recently (127, 128).

ACKNOWLEDGMENTS

The Committee wishes to acknowledge the assistance of B. L. Larson, J. R. Brunner, and S. Nakai in preparation of this manuscript as well as other workers in the field who have commented on proposed changes in nomenclature. Special thanks is extended to J. J. Basch for his electrophoretic comparison of the various preparations of MFGM.

REFERENCES

- 1 Aalund, O., D. Blakeslee, J. E. Butler, J. R. Duncan, M. J. Freeman, R. Jenness, J. M. Kehoe, J.-P. Mach, J. Rapacz, J.-P. Vaerman, and A. J. Winter. 1971. Proposed nomenclature for the immunoglobulins of the domesticated *Bovidae*. *Can. J. Comp. Med.* 35:346.
- 2 Ambrosius, H., R. Asofsky, R. A. Binaghi, D. Capan, L. Hood, E. Orlans, D. S. Rowe, P. Small, G. Torrigiani, and J.-P. Vaerman. 1978. Proposed rules for the designation of immunoglobulins of animal origin. *Bull. World Health Org.* 56:815.
- 3 Aimutis, W. R., and W. N. Eigel. 1982. Identification of λ -casein as plasmin-derived fragments of bovine α_{S1} -casein. *J. Dairy Sci.* 65:175.
- 4 Anderson, M., and T. E. Cawston. 1975. Review of the progress of dairy science. Milk-fat globule membrane. *J. Dairy Res.* 42:459.
- 5 Anderson, M., T. Cawston, and G. C. Cheeseman. 1974. Molecular-weight estimates of milk-fat-globule-membrane protein-sodium dodecyl sulfate complexes by electrophoresis in gradient acrylamide gels. *Biochim. J.* 139:653.
- 6 Anderson, M., and G. C. Cheeseman. 1971. Some aspects of the chemical composition of the milk fat globule membrane during lactation. *J. Dairy Res.* 38:409.
- 7 Andrews, A. T. 1978. The composition, structure and origin of proteose-peptone component 5 of bovine milk. *Eur. J. Biochem.* 90:59.

- 8 Andrews, A. T. 1978. The composition, structure and origin of proteose-peptone component 8F of bovine milk. *Eur. J. J. Biochem.* 90:67.
- 9 Annan, W. D., and W. Manson. 1969. A fractionation of the α_s -casein complex of bovine milk. *J. Dairy Res.* 36:259.
- 10 Aschaffenburg, R. 1963. Milk protein polymorphisms. Page 50 in *Man and cattle*. A. E. Mourant and F. E. Zeuner, ed. Royal Anthropol. Inst., Occas. Paper No. 18, London.
- 11 Aschaffenburg, R. 1968. Reviews of the progress of dairy science. Section G. Genetics. Genetic variants of milk proteins: their breed distribution. *J. Dairy Res.* 35:447.
- 12 Aschaffenburg, R., and J. Drewry. 1957. Genetics of the β -lactoglobulins of cow's milk. *Nature* 180:376.
- 13 Aschaffenburg, R., and J. Drewry. 1957. Improved method for the preparation of crystalline β -lactoglobulin and α -lactalbumin from cow's milk. *Biochem. J.* 65:273.
- 14 Aschaffenburg, R., A. Sen, and M. P. Thompson. 1968. Genetic variants of casein in Indian and African zebu cattle. *Comp. Biochem. Physiol.* 25:177.
- 15 Baldwin, R. L. 1957. Boundary spreading in sedimentation-velocity experiments. 5. Measurement of the diffusion coefficient of bovine albumin by Fujita's equation. *Biochem. J.* 65:503.
- 16 Barman, T. E. 1970. Purification and properties of bovine milk glyco- α -lactalbumin. *Biochim. Biophys. Acta* 214:243.
- 17 Barman, T. E. 1973. The isolation of an α -lactalbumin with three disulfide bonds. *Eur. J. Biochem.* 37:86.
- 18 Basch, J. J., H. M. Farrell, Jr., and R. Greenberg. 1976. Identification of the milk fat globule membrane proteins. I. Isolation and partial characterization of a glycoprotein B. *Biochim. Biophys. Acta* 448:589.
- 19 Basch, J. J., R. Greenberg, and H. M. Farrell, Jr. 1981. Use of preparative gel electrophoresis in the study of the proteins of the milk fat globule membrane. *Fed. Proc.* 40:1660.
- 20 Beeby, R. 1964. The presence of sulphhydryl groups in κ -casein. *Biochim. Biophys. Acta* 82:418.
- 21 Beh, K. J. 1973. Distribution of Brucella antibody among immunoglobulin classes and a low molecular weight antibody fraction in serum and whey of cattle. *Res. Vet. Sci.* 14:381.
- 22 Behrens, P. Q., A. M. Spiekerman, and J. R. Brown. 1975. Structure of human serum albumin. *Fed. Proc.* 34:591.
- 23 Bell, K., K. E. Hopper, and H. A. McKenzie. 1981. Bovine α -lactalbumin C and α_{s1} , β - and κ -caseins of Bali (Banteng) cattle, *Bos (Bibos) javanicus*. *Aust. J. Biol. Sci.* 34:149.
- 24 Bell, K., H. A. McKenzie, W. H. Murphy, and D. C. Shaw. 1970. β -Lactoglobulin (Droughtmaster): A unique protein variant. *Biochim. Biophys. Acta* 214:427.
- 25 Bell, K., H. A. McKenzie, and D. C. Shaw. 1981. Bovine β -lactoglobulin E, F and G of Bali (Banteng) cattle, *Bos (Bibos) javanicus*. *Aust. J. Biol. Sci.* 34:133.
- 26 Benten, Ch. von, W. Floer, and K. Petzoldt. 1976. Untersuchungen zum tierexperimentellen Nachweis von anaphylaktischen Antikörpern des Rindes im passiven Übertragungsversuch unter Verwendung von kolostrumfrei ernährten Kalbern und Lämmern. *Zentralbl. Veterinärmed. B* 23:200.
- 27 Berrens, L., and E. Bleumink. 1965. Spectroscopic evidence for N-glycosidic linkages in atopic allergens. *Int. Arch. Allergy Appl. Immunol.* 28:150.
- 28 Bezkorovainy, A., and D. Grohlich. 1972. Maleylation of bovine colostrum IgG and human serum IgG. *Immunochemistry* 9:137.
- 29 Bhattacharya, S. D., A. K. Roychoudhury, N. K. Sinha, and A. Sen. 1963. Inherited α -lactalbumin and β -lactoglobulin polymorphism in Indian Zebu cattle. Comparison of Zebu and buffalo α -lactalbumins. *Nature* 197:797.
- 30 Bigelow, C. C., and M. Channon. 1976. Hydrophobicities of amino acids and proteins. Page 209 in *Handbook of biochemistry and molecular biology*. Vol. I. G. D. Fasman, ed. CRC Press, Inc., Cleveland, OH.
- 31 Bingham, E. W. 1979. Role of mammary casein kinase in the phosphorylation of milk proteins. *J. Dairy Res.* 46:181.
- 32 Bingham, E. W., and H. M. Farrell, Jr. 1977. Phosphorylation of casein by the lactating mammary gland: A review. *J. Dairy Sci.* 60:1199.
- 33 Bingham, E. W., H. M. Farrell, Jr., and R. Carroll. 1972. Properties of dephosphorylated α_{s1} -casein. Precipitation by calcium ions and micelle formation. *Biochemistry* 11:2450.
- 34 Brandon, M. R., D. L. Watson, and A. K. Lascelles. 1971. The mechanism of transfer immunoglobulin into mammary secretion of cows. *Aust. J. Exp. Biol. Med. Sci.* 49:613.
- 35 Braunitzer, G., and R. Chen. 1972. Die Spaltung des β -Lactoglobulins AB mit Bromcyan. *Hoppe-Seyler's Z. Physiol. Chem.* 353:674.
- 36 Braunitzer, G., R. Chen, B. Schrank, and A. Stangl. 1972. Automatische Sequenzanalyse eines Proteins (β -Lactoglobulin AB). *Hoppe-Seyler's Z. Physiol. Chem.* 353:832.
- 37 Brew, K., F. J. Castellino, T. C. Vanaman, and R. L. Hill. 1970. The complete amino acid sequence of bovine α -lactalbumin. *J. Biol. Chem.* 245:4570.
- 38 Brew, K., and R. L. Hill. 1970. The isolation and characterization of the tryptic, chymotryptic, peptic, and cyanogen bromide peptides from bovine α -lactalbumin. *J. Biol. Chem.* 245:4559.
- 39 Brew, K., and R. L. Hill. 1975. Lactose biosynthesis. *Rev. Physiol. Biochem. Pharmacol.* 72:105.
- 40 Brignon, G., and B. Ribadeau-Dumas. 1973. Localisation dans la chaîne peptidique de la β lactoglobuline bovine de la substitution Glu/Gln différenciant les variants génétiques B et D. *FEBS Lett.* 33:73.
- 41 Brignon, G., B. Ribadeau-Dumas, and J.-C.

- Mercier. 1976. Primiers elements de structure primaire des caseines α_{s2} bovines. *FEBS Lett.* 71:111.
- 42 Brignon, G., B. Ribadeau-Dumas, J.-C. Mercier, J.-P. Pelissier, and B. C. Das. 1977. Complete amino acid sequence of bovine α_{s2} -casein. *FEBS Lett.* 76:274.
 - 43 Brown, J. R. 1975. Structure of bovine serum albumin. 1975. *Fed. Proc.* 34:591.
 - 44 Brown, J. R. 1977. Structure and evolution of serum albumin. In *Albumin Structure Biosynthesis; Function*. Proc. FEBS Meet. 50:1.
 - 45 Brunner, J. R. 1974. Physical equilibria in milk: the lipid phase. Page 474 in *Fundamentals of dairy chemistry*. B. H. Webb, A. H. Johnson, and J. A. Alford, ed. Avi Publ. Co., Inc., Westport, CT.
 - 46 Brunner, J. R. 1977. Milk proteins. Page 175 in *Food proteins*. J. R. Whitaker and S. R. Tanenbaum, ed. Avi Publ. Co., Inc., Westport, CT.
 - 47 Brunner, J. R. 1981. Cow milk proteins: Twenty-five years of progress. *J. Dairy Sci.* 64:1038.
 - 48 Brunner, J. R., C. A. Ernstrom, R. A. Hollis, B. L. Larson, R. McL. Whitney, and C. A. Zittle. 1960. Nomenclature of the proteins of bovine milk - first revision. *J. Dairy Sci.* 43:901.
 - 49 Butler, J. E. 1969. Bovine immunoglobulins: A review. *J. Dairy Sci.* 52:1895.
 - 50 Butler, J. E. 1971. Physicochemical and immunochemical studies of bovine IgA and glycoprotein-a. *Biochim. Biophys. Acta* 251:435.
 - 51 Butler, J. E. 1973. The occurrence of immunoglobulin fragments, two types of lactoferrin and a lactoferrin-IgG₂ complex in bovine colostrum and milk whey. *Biochim. Biophys. Acta* 295:341.
 - 52 Butler, J. E. 1974. Immunoglobulins of the mammary secretions. Page 217 in *Lactation, a comprehensive treatise*. Vol. III. B. L. Larson and V. R. Smith, ed. Academic Press, New York, NY.
 - 53 Butler, J. E. 1979. Immunologic aspects of breast feeding, anti-infections activity of breast milk. Page 255 in *Human lactation. Seminars in perinatology*. Vol. 3. T. K. Oliver and T. H. Kirschbaum, ed.
 - 54 Butler, J. E. 1981. A concept of humoral immunity among ruminants and an approach to its investigation. Page 10 in *The ruminant immune system*. J. E. Butler, ed. Plenum Press, New York, NY.
 - 55 Butler, J. E. 1983. Bovine immunoglobulins: An augmented review. *Vet. Immunol. Immunopathol.* 4:43.
 - 56 Butler, J. E., E. J. Coulson, and M. L. Groves. 1968. Identification of glycoprotein-a as a probable fragment of bovine IgA. *Fed. Proc.* 27:617.
 - 57 Butler, J. E., C. A. Kiddy, C. S. Pierce, and C. A. Rock. 1972. Quantitative changes associated with calving in the levels of bovine immunoglobulins in selected body fluids. I. Changes in the levels of IgA, IgG1 and total protein. *Can. J. Comp. Med.* 36:234.
 - 58 Butler, J. E., P. L. McGivern, L. A. Cantarero, and L. Peterson. 1980. Application of the amplified enzyme-linked immunosorbent assay: Comparative quantitation of bovine serum IgG1, IgG2, IgA and IgM antibodies. *Am. J. Vet. Res.* 41:1479.
 - 59 Butler, J. E., L. Peterson, and P. McGivern. 1980. A reliable method for the preparation of bovine secretory immunoglobulin A (SIgA) which circumvents problems posed by IgG1 dimers in colostrum. *Mol. Immunol.* 17:757.
 - 60 Butler, J. E., A. J. Winter, and G. G. Wagner. 1971. Symposium: Bovine immune systems. *J. Dairy Sci.* 54:1309.
 - 61 Ceppellini, R., S. Dray, G. Edelman, J. Fahey, F. Franek, E. Franklin, H. C. Goodman, P. Grabar, A. E. Gurvich, J. F. Heremans, H. Isliker, F. Karush, E. Press, and Z. Truka. 1964. Nomenclature for human immunoglobulins. *Bull. World Health Org.* 30:447.
 - 62 Coulson, E. J., and H. Stevens. 1950. The serological relationship of bovine whey albumin to serum albumin. *J. Biol. Chem.* 187:355.
 - 63 Dayhoff, M. O. 1976. Page 267 in *Atlas of protein sequence and structure*. M. O. Dayhoff, ed. Vol. 5. Suppl. 2. Natl. Biomed. Res. Found., Washington, DC.
 - 64 Delfour, A., J. Jollés, C. Alais, and P. Jollés. 1965. Caseino-glycopeptides. Characterization of a methionine residue and of the N-terminal sequence. *Biochem. Biophys. Res. Comm.* 19:452.
 - 65 Doi, H., F. Ibuki, and M. Kanamori. 1979. Heterogeneity of reduced bovine κ -casein. *J. Dairy Sci.* 62:195.
 - 66 Doi, H., N. Kawaguchi, F. Ibuki, and M. Kanamori. 1979. Minor components of reduced κ -casein. *J. Nutr. Vitaminol.* 25:95.
 - 67 Doi, H., H. Kobatake, F. Ibuki, and M. Kanamori. 1980. Attachment sites of carbohydrate portions to peptide chain of κ -casein from bovine colostrum. *Agric. Biol. Chem.* 44:2605.
 - 68 Duncan, J. R., B. N. Wilkie, F. Hiestand, and A. J. Winter. 1972. The serum and secretory immunoglobulins of cattle: Characterization and quantitation. *J. Immunol.* 108:965.
 - 69 Ebner, K. E., W. L. Denton, and U. Brodbeck. 1966. The substitution of α -lactalbumin for the B protein of lactose synthetase. *Biochem. Biophys. Res. Comm.* 24:232.
 - 70 Ebner, K. E., and F. Schanbacher. 1974. Biochemistry of lactose and related carbohydrates. Page 77 in *Lactation: A comprehensive Treatise*. Vol. II. B. L. Larson and V. R. Smith, ed. Academic Press, New York, NY.
 - 71 Eigel, W. N. 1977. Formation of γ_1 -A², γ_2 -A² and γ_3 -A caseins by in vitro proteolysis of β -casein A² with bovine plasmin. *Int. J. Biochem.* 8:187.
 - 72 Eigel, W. N. 1981. Identification of proteose-peptone component 5 as a plasmin-derived fragment of bovine β -casein. *Int. J. Biochem.* 13:1081.
 - 73 Eigel, W. N., C. J. Hofmann, B.A.K. Chibber, J. M. Tomich, T. W. Keenen, and E. T. Mertz. 1979. Plasmin-mediated proteolysis of casein in bovine milk. *Proc. Natl. Acad. Sci.* 76:2244.

- 74 Eigel, W. N., and T. W. Keenan. 1979. Identification of proteose peptone component 8-slow as a plasmin-derived fragment of bovine β -casein. *Int. J. Biochem.* 10:529.
- 75 Ehrenpreis, S., P. H. Maurer, and J. Sri Ram. 1957. Modified bovine serum albumin. I. Preparation and physicochemical studies of some derivatives. *Arch. Biochem. Biophys.* 67:178.
- 76 El-Negoumy, A. M. 1973. Separation of lambda casein and some of its properties. *J. Dairy Sci.* 56:1486.
- 77 Evans, M.T.A., L. Irons, and M. Jones. 1971. Physicochemical properties of β -casein and some carboxyacyl derivatives. *Biochim. Biophys. Acta* 229:411.
- 78 Feinstein, A., and M. J. Hobart. 1969. Structural relationship and complement fixing activity of sheep and other ruminant immunoglobulin G subclasses. *Nature* 223:950.
- 79 Fey, H., H. Pfister, J. Messerli, N. Sturzenegger, and F. Grolimund. 1976. Methods of isolation purification and quantitation of the bovine immunoglobulins. A technical review. *Zentrabl. Veterinaarmed. B* 23:269.
- 80 Fiat, A.-M., C. Alais, and P. Jollés. 1972. The amino-acid and carbohydrate sequences of a short glycopeptide isolated from bovine κ -casein. *Eur. J. Biochem.* 27:408.
- 81 Fournet, B., A.-M. Fiat, C. Alais, and P. Jollés. 1979. Cow κ -casein: Structure of the carbohydrate portion. *Biochim. Biophys. Acta* 576:339.
- 82 Fournet, B., A.-M. Fiat, J. Montreuil, and P. Jollés. 1975. The sugar part of κ -caseins from cow milk and colostrum and its microheterogeneity. *Biochimie* 57:161.
- 83 Fox, P. F. 1982. Heat-induced coagulation of milk. Page 189 in *Developments in dairy chemistry*. P. F. Fox, ed. Appl. Sci. Publ., LTD, London.
- 84 Fox, P. F., and D. M. Mulvihill. 1982. Milk proteins: molecular, colloidal and functional properties. *J. Dairy Res.* 49:679.
- 85 Frank, V. G., and G. Braunitzer. 1967. Zur Primärstruktur der β -Lactoglobuline. *Hoppe-Seyler's Z. Physiol. Chem.* 348:1691.
- 86 Frank, V. G., and G. Braunitzer. 1968. Gewinnung aller tryptischen Peptide von β -Lactoglobulin A and B. *Hoppe-Seyler's Z. Physiol. Chem.* 349:1456.
- 87 Franke, W. W., H. W. Heide, C. Gruno, S. Winter, C. Freudenstein, E. Schmid, E.-D. Jarasch, and T. W. Keenan. 1981. Antibodies to the major insoluble milk fat globule membrane protein: Specific location in apical regions of lactating cells. *J. Cell Biol.* 89:485.
- 88 Fredriksson, S. 1975. Scanning isoelectric focusing in small density-gradient columns. IV. The use of deuterium oxide for preparing the density gradient and its effects on isoelectric points of proteins. *J. Chromatogr.* 108:153.
- 89 Freudenstein, C., T. W. Keenan, W. N. Eigel, M. Sasaki, J. Stadler, and W. W. Franke. 1979. Preparation and characterization of the inner coat material associated with fat globule membranes from bovine and human milk. *Exp. Cell Res.* 118:277.
- 90 Gally, J. A. 1973. Structure of immunoglobulins. Page 162 in *The antigens*. Vol. I. M. Sela, ed. Academic Press, New York, NY.
- 91 Garnier, J. 1963. Étude de l'action de la pression sur la caséine par la méthode titrimétrique a pH constant. Comparaison avec d'autres méthodes. *Ann. Biol. Anim. Biochem. Biophys.* 3:71.
- 92 Goodger, B. V. 1971. A low molecular weight protein in bovine serum with similar electrophoretic mobility to IgG₂. *Res. Vet. Sci.* 12:465.
- 93 Gordon, W. G. 1971. α -Lactalbumin. Page 331 in *Milk proteins*. Vol. II. H. A. McKenzie, ed. Academic Press, New York, NY.
- 94 Gordon, W. G., R. Aschaffenburg, A. Sen, and S. K. Ghosh. 1968. Amino acid composition of several α -lactalbumins. *J. Dairy Sci.* 51:947.
- 95 Gordon, W. G., M. L. Groves, R. Greenberg, S. B. Jones, E. B. Kalan, R. F. Peterson, and R. E. Townend. 1972. Probable identification of γ -, TS-, R- and S-caseins as fragments of β -casein. *J. Dairy Sci.* 55:261.
- 96 Gough, P., R. Jenness, and R. K. Anderson. 1966. Characterization of bovine immunoglobulins. *J. Dairy Sci.* 49:718.
- 97 Grosclaude, F., P. Joudrier, and M.-F. Mahé. 1978. Polymorphisme de la caséine α_{S2} -bovine: étroite liaison du locus α_{S2} -Cn avec les loci α_{S1} -Cn, β -Cn et κ -Cn; mise en évidence d'une deletion dans le variant α_{S2} -Cn D. *Ann. Genet. Sel. Anim.* 10:313.
- 98 Grosclaude, F., P. Joudrier, and M.-F. Mahé. 1979. A genetic and biochemical analysis of a polymorphism of bovine α_{S2} -casein. *J. Dairy Res.* 46:211.
- 99 Grosclaude, F., M.-F. Mahé, and J.-C. Mercier. 1974. Comparaison du polymorphisme genetique des lactoprotéines du zebu et des bovines. *Ann. Génét. Sel. Anim.* 6:305.
- 100 Grosclaude, F., M.-F. Mahé, J.-C. Mercier, J. Bonnemaire, and J. H. Teissier. 1976. Polymorphisme des lactoprotéines de bovines Népalaïs. I. Mise en évidence, chez le Yak, et caractérisation biochimique de deux nouveaux variants: β -lactoglobuline D (yak) et caséine α_{S1} -E. *Ann. Génét. Sel. Anim.* 8:461.
- 101 Grosclaude, F., M.-F. Mahé, J.-C. Mercier, J. Bonnemaire, and J. H. Teissier. 1976. Polymorphisme de lactoprotéines de bovines Népalaïs. II. Polymorphisme des caséines (α_S -mineures); Le locus α_{S2} -Cn est-il lié aux loci α_{S1} -Cn, β -Cn, in κ -Cn? *Ann. Génét. Sél. Anim.* 8:481.
- 102 Grosclaude, F., M.-F. Mahé, J.-C. Mercier, and B. Ribadeau-Dumas. 1972. Caractérisation des variants génétiques des caséines α_{S1} et β bovines. *Eur. J. Biochem.* 26:328.
- 103 Grosclaude, F., M.-F. Mahé, and B. Rabadeau-Dumas. 1973. Structure primaire de la caséine α_{S1} , et de la caséine β bovine. *Eur. J. Biochem.* 40:323.
- 104 Grosclaude, F., M.-F. Mahé, and G.-F. Voglino. 1974. Le variant β E et le code de phosphorylation des caséines bovines. *FEBS Lett.* 45:3.

- 105 Groves, M. L. 1960. The isolation of a red protein from milk. *J. Am. Chem. Soc.* 82:3345.
- 106 Groves, M. L. 1971. Minor milk proteins and enzymes. Page 367 in *Milk proteins*. Vol. II. H. A. McKenzie, ed. Academic Press, New York, NY.
- 107 Groves, M. L., J. J. Basch, and W. G. Gordon. 1963. Isolation, characterization and amino acid composition of a new crystalline protein, lactollin, from milk. *Biochemistry* 2:814.
- 108 Groves, M. L., and W. G. Gordon. 1967. Isolation of a new glycoprotein-a and a γ G-globulin from individual cow milks. *Biochemistry* 6:2388.
- 109 Groves, M. L., W. G. Gordon, E. B. Kalan, and S. B. Jones. 1972. Composition of bovine γ -caseins A¹ and A² and further evidence for a relationship in biosynthesis of γ - and β -caseins. *J. Dairy Sci.* 55:1041.
- 110 Groves, M. L., W. G. Gordon, E. B. Kalan, and S. B. Jones. 1973. TS-A², TS-B, R- and S-caseins: Their isolation, composition and relationship to the β - and γ -casein polymorphs A² and B. *J. Dairy Sci.* 56:558.
- 111 Groves, M. L., and R. Greenberg. 1977. Bovine homologue of β_2 -microglobulin isolated from milk. *Biochem. Biophys. Res. Comm.* 77:320.
- 112 Groves, M. L., and R. Greenberg. 1982. Complete amino acid sequence of bovine β_2 -microglobulin. *J. Biol. Chem.* 257:2619.
- 113 Guidry, A. J., J. E. Butler, R. E. Pearson, and B. T. Weinland. 1980. IgA, IgG1, IgG2, IgM, and BSA in serum and mammary secretion throughout lactation. *Vet. Immunol. Immunopathol.* 1:329.
- 114 Halbeek, H. van, L. Dorland, J.F.G. Vliegthart, A.-M. Fiat, and P. Jollés. 1981. Structural characterization of a novel acidic oligosaccharide unit derived from cow colostrum κ -casein. *FEBS Lett.* 133:45.
- 115 Hammer, D. K., B. Kickhöfen, and T. Schmid. 1971. Detection of homocytotropic antibody associated with a unique immunoglobulin class in the bovine species. *Eur. J. Immunol.* 1:249.
- 116 Herald, C. T., and J. R. Brunner. 1957. The fat-globule membrane of normal cow's milk. I. The isolation and characteristics of two membrane-protein fractions. *J. Dairy Sci.* 40:948.
- 117 Hill, R. L., and K. Brew. 1975. Lactose synthetase. *Adv. Enzymol. Relat. Areas Mol. Biol.* 43:411.
- 118 Hiraoka, Y., T. Segawa, K. Kuwajima, S. Sugai, and N. Murai. 1980. α -Lactalbumin: A calcium metalloprotein. *Biochem. Biophys. Res. Comm.* 95:1098.
- 119 Ho, C., and D. F. Waugh. 1965. Interactions of bovine α_s -casein with small ions. *J. Am. Chem. Soc.* 87:110.
- 120 Hoagland, P. D., M. P. Thompson, and E. B. Kalan. 1971. Amino acid composition of α_{s3} , α_{s4} , and α_{s5} -caseins. *J. Dairy Sci.* 54:1103.
- 121 Hofmann, C. J., T. W. Keenan, and W. N. Eigel. 1979. Association of plasminogen with bovine milk fat globule membrane. *Int. J. Biochem.* 10:909.
- 122 Honacki, J. H., K. E. Kinman, and J. W. Koepl. 1982. *Mammal species of the world*. Assoc. Systematics Collections and Allen Press, Inc., Lawrence, KS.
- 123 Hood, L., W. R. Gray, B. G. Sanders, and W. J. Dreyer. 1967. Light chain evolution. *Cold Spring Harbor Symp. Quant. Biol.* 32:133.
- 124 Hopper, K. E., and H. A. McKenzie. 1973. Minor components of bovine α -lactalbumin A and B. *Biochim. Biophys. Acta* 295:352.
- 125 Jarasch, E.-D., G. Bruder, T. W. Keenan, and W. W. Franke. 1977. Redox constituents in milk fat globule membranes and rough endoplasmic reticulum from lactating mammary gland. *J. Cell Biol.* 73:223.
- 126 Jenness, R. 1970. Protein composition of milk. Page 17 in *Milk proteins*. Vol. I. H. A. McKenzie, ed. Academic Press, New York, NY.
- 127 Jenness, R. 1979. Comparative aspects of milk proteins. *J. Dairy Res.* 46:197.
- 128 Jenness, R. 1982. Interspecies comparison of milk proteins. Page 87 in *Developments in dairy chemistry*. Vol. I. P. F. Fox, ed. Appl. Sci. Publ., London.
- 129 Jenness, R., B. L. Larson, T. L. McMeekin, A. M. Swanson, C. H. Whitnah, and R. McL. Whitney. 1956. Nomenclature of the proteins of bovine milk. *J. Dairy Sci.* 39:536.
- 130 Jollés, P. 1975. Structural aspects of the milk clotting process. Comparative features with the blood clotting process. *Mol. Cell Biochem.* 7:73.
- 131 Jollés, J., C. Alais, and P. Jollés. 1968. The tryptic peptide with the rennin-sensitive linkage of cow's κ -casein. *Biochim. Biophys. Acta* 168:591.
- 132 Jollés, J., A.-M. Fiat, C. Alais, and P. Jollés. 1973. Comparative study of cow and sheep κ -caseinoglycopeptides: Determination of the N-terminal sequences with a sequencer and the location of the sugars. *FEBS Lett.* 30:173.
- 133 Jollés, P., M.-H. Loucheux-Lefebvre, and A. Henschen. 1978. Structural relatedness of κ -casein and fibrinogen γ -chain. *J. Mol. Evol.* 11:271.
- 134 Jollés, J., F. Schoentgen, C. Alais, A.-M. Fiat, and P. Jollés. 1972. Studies on the primary structure of cow κ -casein. Structural features of para- κ -casein; N-terminal sequence of κ -caseinoglycopeptide studied with a sequencer. *Helv. Chim. Acta* 55:2872.
- 135 Joniau, M., J. Bloemmen, and R. Lontie. 1970. The reaction of 4-(p-sulphophenylazo)-2-mercuriphenol with the thiol groups of proteins. *Biochim. Biophys. Acta* 214:468.
- 136 Jones, E. A. 1977. Synthesis and secretion of milk sugars. *Symp. Zool. Soc. London* 41:77.
- 137 Jones, E. A. 1978. Lactose biosynthesis. Page 371 in *Lactation: a comprehensive treatise*. Vol. IV. B. L. Larson, ed. Academic Press, New York, NY.
- 138 Josephson, R. V., 1972. Isoelectric focusing of bovine milk caseins. *J. Dairy Sci.*, 55:1535.
- 139 Josephson, R. V., E. M. Mikolajcik, and D. P. Sinha. 1972. Gel isoelectric focusing of selected bovine immunoglobulins. *J. Dairy Sci.* 55:1508.
- 140 Kaldor, G., A. Saifer, and F. Westley. 1962. Physicochemical studies of ultraviolet-irradiated bovine serum albumins. *Arch. Biochem. Biophys.* 99:275.

- 141 Kaminogawa, S., H. Mizobuchi, and K. Yamauchi. 1972. Comparison of bovine milk protease with plasmin. *Agric. Biol. Chem.* 36:2163.
- 142 Kanamori, M., N. Kawaguchi, F. Ibuki, and H. Doi. 1980. Attachment sites of carbohydrate moieties to peptide chain of bovine κ -casein from normal milk. *Agric. Biol. Chem.* 44:1855.
- 143 Kanno, C., D. B. Emmons, V. R. Harwalkar, and J. A. Elliott. 1976. Purification and characterization of the agglutinating factor for lactic Streptococci from bovine milk: IgM-Immunoglobulin. *J. Dairy Sci.* 59:2036.
- 144 Kanno, C., M. Shimizu, and K. Yamauchi. 1975. Isolation and physicochemical properties of a soluble glycoprotein fraction of milk fat globule membrane. *Agric. Biol. Chem.* 39:1835.
- 145 Kanno, C., and K. Yamauchi. 1979. Relationship of soluble glycoprotein of milk fat globule membrane to component -3, -5, and -8 fractions of proteose-peptone. *Agric. Biol. Chem.* 43:2105.
- 146 Keenan, T. W., D. E. Olson, and H. H. Mollenhauer. 1971. Origin of the milk fat globule membrane. *J. Dairy Sci.* 54:295.
- 147 Keenan, T. W., D. J. Morré, D. E. Olson, W. N. Yunghans, and S. Patton. 1970. Biochemical and morphological comparison of plasma membrane and milk fat globule membrane from bovine mammary gland. *J. Cell Biol.* 44:80.
- 148 Kickhöfen, B., D. K. Hammer, and D. Scheel. 1968. Isolation and characterization of γ G type immunoglobulins from bovine serum and colostrum. *Hoppe-Seyler's Z. Physiol. Chem.* 349:1755.
- 149 Kiddy, C. A. 1975. Gel electrophoresis in vertical polyacrylamide beds. Procedures I and II. Page 14 in *Methods of gel electrophoresis of milk proteins*. H. E. Swaisgood, B. L. Larson, E. B. Kalan, J. R. Brunner, C. V. Morr, and P.M.T. Hansen, ed. Am. Dairy Sci. Assoc., Champaign, IL.
- 150 King, N. 1955. The milk fat globule membrane and some associated phenomena. Tech. Commun. No. 2, Commonw. Bur. Dairy Sci., Commonw. Agric. Bur., Farnham Royal, Bucks, England.
- 151 Kitchen, B. J. 1977. Fractionation and characterization of the membranes from bovine milk fat globules. *J. Dairy Res.* 44:469.
- 152 Kobayashi, K., J.-P. Vaerman, H. Bazin, A.-M. Lebacqz-Verheyden, and J. F. Heremans. 1973. Identification of J-chain in polymeric immunoglobulins from a variety of species by cross-reaction with rabbit antisera to human J-chain. *J. Immunol.* 111:1590.
- 153 Kobylka, D., and K. L. Carraway. 1972. Proteins and glycoproteins of the milk fat globule membrane. *Biochim. Biophys. Acta* 288:282.
- 154 Kobylka, D., and K. L. Carraway. 1973. Proteolytic digestion of proteins of the milk fat globule membrane. *Biochim. Biophys. Acta* 307:133.
- 155 Komar, R., and T.K.S. Mukkur. 1975. Isolation and characterization of J-chain from bovine colostrum immunoglobulin. *Can. J. Biochem.* 53:943.
- 156 Kronman, M. J., and R. E. Andreotti. 1964. Inter- and intramolecular interactions of α -lactalbumin. I. The apparent heterogeneity at acid pH. *Biochemistry* 3:1145.
- 157 Kronman, M. J., R. E. Andreotti, and R. Vitols. 1964. Inter- and intramolecular interactions of α -lactalbumin. II. Aggregation reactions at acid pH. *Biochemistry* 3:1152.
- 158 Kumar, S., and E. M. Mikolajcik. 1973. Selected physico-chemical characteristics of bovine colostrum immunoglobulin. *J. Dairy Sci.* 56:255.
- 159 Kumosinski, T. F., E. M. Brown, and M. L. Groves. 1981. Solution physico-chemical properties of bovine B₂-microglobulin. *J. Biol. Chem.* 256:10949.
- 160 Labib, R. S., N. J. Calvanico, and T. B. Tomasi. 1976. Bovine secretory component. Isolation, molecular size and shape, composition, and NH₂-terminal amino acid sequence. *J. Biol. Chem.* 251:1969.
- 161 Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680.
- 162 Lancet, D., P. Parham, and J. L. Strominger. 1979. Heavy chain of HLA-A and HLA-B antigens is conformationally labile: A possible role for B₂-microglobulin. *Proc. Natl. Acad. Sci.* 76:3844.
- 163 Larson, B. L. 1979. Biosynthesis and secretion of milk proteins: A review. *J. Dairy Res.* 46:161.
- 164 Lisowski, J., M. Janusz, B. Tyran, A. Morawiecki, S. Golab, and H. Białkowska. 1975. Immunoglobulins of colostrum - IV: Comparative studies of bovine serum and colostrum IgG1 and IgG2. *Immunochemistry* 12:167.
- 165 Long, J., Q. Van Winkel, and I. A. Gould. 1958. Isolation and identification of λ -casein. *J. Dairy Sci.* 41:317.
- 166 MacDonald, C. A., and M.A.W. Thomas. 1970. The rennin-sensitive bond of bovine κ -casein. *Biochim. Biophys. Acta* 207:139.
- 167 Mach, J.-P. 1970. In vitro combination of human and bovine free secretory component with IgA of various species. *Nature* 228:1278.
- 168 Mach, J.-P., and J.-J. Pahud. 1971. Secretory IgA, a major immunoglobulin in most bovine external secretions. *J. Immunol.* 106:552.
- 169 Mach, J.-P., J.-J. Pahud, and H. Isliker. 1969. IgA with "Secretory Piece" in bovine colostrum and saliva. *Nature* 223:952.
- 170 Mackenzie, D.D.S., and A. K. Lascelles. 1968. The transfer of iodine-131 labelled immunoglobulins and serum albumin from blood into milk of lactating ewes. *Aust. J. Exp. Biol. Med. Sci.* 46:285.
- 171 Mackinlay, A. G., R. J. Hill, and R. G. Wake. 1966. The action of rennin on κ -casein. The heterogeneity and origin of the insoluble products. *Biochim. Biophys. Acta* 115:103.
- 172 Mackinlay, A. G., and R. G. Wake. 1965. Fractionation of S-carboxymethyl- κ -casein and characterization of the components. *Biochim. Biophys. Acta* 104:167.
- 173 Mangino, M. E., and J. R. Brunner. 1975. Molecular weight profile of fat globule membrane

- proteins. *J. Dairy Sci.* 58:313.
- 174 Manson, W., W. D. Annan, and G. K. Barnes. 1976. α_{S0} -casein: Its preparation and characterization. *J. Dairy Res.* 43:133.
- 175 Manson, W., T. Carolan, and W. D. Annan. 1977. Bovine α_{S0} casein; a phosphorylated homologue of α_{S1} casein. *Eur. J. Biochem.* 78:411.
- 176 Marchalonis, J. J., and J. K. Weltman. 1971. Relatedness among proteins: A new method of estimation and its application to immunoglobulins. *Comp. Biochem. Physiol.* 38B:609.
- 177 Martens, C. L. 1978. A glycoprotein of the milk fat globule membrane. Ph.D. thesis, Univ. Iowa.
- 178 Mather, I. H., and T. W. Keenan. 1975. Studies on the structure of the milk fat globule membrane. *J. Membr. Biol.* 21:65.
- 179 Mather, I. H., K. Weber, and T. W. Keenan. 1977. Membranes of mammary gland. XII. Loosely associated proteins and compositional heterogeneity of bovine milk fat globule membrane. *J. Dairy Sci.* 60:394.
- 180 McKenzie, H. A. 1971. β -lactoglobulin. Page 257 in *Milk proteins*. Vol. II. H. A. McKenzie, ed. Academic Press, New York, NY.
- 181 McKenzie, R. M., and B. L. Larson. 1978. Purification and partial characterization of a unique group of phosphoproteins from rat milk whey. *J. Dairy Sci.* 61:723.
- 182 McKenzie, H. A., G. B. Ralston, and D. C. Shaw. 1972. Location of sulphhydryl and disulfide groups in bovine β -lactoglobulins and effects of urea. *Biochemistry* 11:4539.
- 183 Mehta, P. D., M. Reichlin, and T. B. Tomasi, Jr. 1972. Comparative studies of vertebrate immunoglobulins. *J. Immunol.* 109:1272.
- 184 Meloun, B., L. Morávek, and V. Kostka. 1975. Complete amino acid sequence of human serum albumin. *FEBS Lett.* 58:134.
- 185 Mephams, T. B. 1977. Synthesis and secretion of milk proteins. *Symp. Zool. Soc. London* 41: 57.
- 186 Mercier, J.-C., G. Brignon, and B. Ribadeau-Dumas. 1973. Structure primaire de la caséine κ B bovine. Séquence complète. *Eur. J. Biochem.* 35:222.
- 187 Mercier, J.-C., F. Grosclaude, and B. Ribadeau-Dumas. 1971. Structure primaire de la caséine α_{S1} -bovine. *Eur. J. Biochem.* 23:41.
- 188 Mukkur, T.K.S., and A. Froese. 1971. Isolation and characterization of IgM from bovine colostrum whey. *Immunochemistry* 8:257.
- 189 Neelin, J. M. 1964. Variants of κ -casein revealed by improved starch gel electrophoresis. *J. Dairy Sci.* 47:506.
- 190 Neville, D. M., and H. Glossmann. 1971. Plasma membrane protein subunit composition. A comparative study by discontinuous electrophoresis in sodium dodecyl sulfate. *J. Biol. Chem.* 246:6335.
- 191 Nezlin, R. S. 1972. Recommendations on the nomenclature of immunoglobulins. (Suggestions of the Subcommittee for Human Immunoglobulins under the Nomenclature Committee of the International Union of Immunological Societies.) *Mol. Biol.* 6:639.
- 192 Nielsen, K. H. 1977. Bovine reaginic antibody. III. Cross-reaction of antihuman IgE and anti-bovine reaginic immunoglobulin antisera with sera from several species of mammals. *Can. J. Comp. Med.* 41:345.
- 193 Nielsen, C. S., and O. J. Bjerrum. 1977. Crossed immunoelectrophoresis of bovine milk fat globule membrane protein solubilized with non-ionic detergent. *Biochim. Biophys. Acta* 466:496.
- 194 Nobel, R. W., Jr., and D. F. Waugh. 1965. Casein micelles. Formation and structure. I. *J. Am. Chem. Soc.* 87:2236.
- 195 Payens, T.A.J., and B. W. van Markwijk. 1963. Some features of the association of β -casein. *Biochim. Biophys. Acta* 71:517.
- 196 Patton, S., and T. W. Keenan. 1975. The milk fat globule membrane. *Biochim. Biophys. Acta* 415:273.
- 197 Pearce, R. J. 1980. Heat-stable components in the Aschaffenburg and Drewry total albumin fraction from bovine milk. *N.Z. J. Dairy Sci. Technol.* 15:13.
- 198 Peters, T., Jr., and R. G. Reed. 1977. Serum Albumin: Conformation and active sites. *Proc. FEBS Mtg.* 50:1.
- 199 Peterson, R. F. 1963. High resolution of milk proteins obtained by gel electrophoresis. *J. Dairy Sci.* 46:1136.
- 200 Peterson, P. A., B. A. Cunningham, I. Berggard, and G. M. Edelman. 1972. β_2 -Microglobulin-A free immunoglobulin domain. *Proc. Natl. Acad. Sci.* 69:1697.
- 201 Phillips, D. R., and P. P. Agin. 1976. Determination of nonequivalent inter and intramolecular disulfide bonds in platelet plasma membrane glycoproteins by two dimensional electrophoresis. *Fed. Proc.* 35:1379.
- 202 Piletz, J. E., M. Heinlen, and R. E. Ganschow. 1981. Biochemical characterization of a novel whey protein from murine milk. *J. Biol. Chem.* 256:11509.
- 203 Polis, B. D., H. W. Shmukler, and J. H. Custer. 1950. Isolation of a crystalline albumin from milk. *J. Biol. Chem.* 187:349.
- 204 Polzhofer, K. P. 1972. Synthese eines labempfindlichen Pentadeca peptids aus kuh- κ -Casein. *Tetrahedron* 28:855.
- 205 Porter, P. 1971. Immunoglobulin IgA in bovine mammary secretions and serum of the neonatal calf. *Biochim. Biophys. Acta* 236:664.
- 206 Porter, P., and D. E. Noakes. 1970. Immunoglobulin IgA in bovine serum and external secretions. *Biochem. Biophys. Acta* 214:107.
- 207 Préaux, G., G. Braunitzer, B. Schrank, and A. Stangl. 1979. The amino acid sequence of goat β -lactoglobulin. *Hoppe-Seyler's Z. Physiol. Chem.* 360:1595.
- 208 Pringnitz, D. J. 1980. Quantitative studies on the origin and distribution of bovine associated mucoprotein in the cow. Ph.D. Thesis, Univ. Iowa.
- 209 Pringnitz, D. J., and J. E. Butler. 1981. The response of rabbits to the subclass specific epitopes of bovine IgG₁ and IgG₂. *Vet. Im-*

- munol. Immunopathol. 2:353.
- 210 Pujolle, J., B. Ribadeau-Dumas, J. Garnier, and R. Pion. 1966. A study of κ -casein components. *Biochem. Biophys. Res. Comm.* 25:285.
 - 211 Quarfoth, G. J., and R. Jenness. 1975. Isolation, composition and functional properties of α -lactalbumins from several species. *Biochim. Biophys. Acta* 379:476.
 - 212 Radl, J., F. Klein, P. van den Berg, A. M. de Bruyn, and W. Hijmans. 1971. Binding of secretory piece to polymeric IgA and IgM paraproteins in vitro. *Immunology* 20:843.
 - 213 Radola, B. J. 1973. Isoelectric focusing in layers of granulated gels. I. Thin-layer isoelectric focusing of proteins. *Biochim. Biophys. Acta* 295:412.
 - 214 Reed, R. G., F. W. Putnam and T. Peters, Jr. 1980. Sequence of residues 400-403 of bovine serum albumin. *Biochem. J.* 191:867.
 - 215 Reynolds, H. Y., and J. S. Johnson. 1971. Structural units of canine serum and secretory immunoglobulin A. *Biochemistry* 10:2821.
 - 216 Ribadeau-Dumas, B., G. Brignon, F. Grosclaude, and J.-C. Mercier. 1972. Structure primaire de la casein β bovine. *Eur. J. Biochem.* 25:505.
 - 217 Richardson, B. C., and L. K. Creamer. 1974. Comparative micelle structure. III. The isolation and chemical characterization of caprine β_1 -casein and β_2 -casein. *Biochim. Biophys. Acta* 365:133.
 - 218 Rose, D., J. R. Brunner, E. B. Kalan, B. L. Larson, P. Melnychyn, H. E. Swaisgood, and D. F. Waugh. 1970. Nomenclature of the proteins of cow's milk: Third revision. *J. Dairy Sci.* 53:1.
 - 219 Saito, T., T. Itoh, and S. Adachi. 1981. The chemical structure of a tetrasaccharide containing N-acetylglucosamine obtained from bovine colostrum κ -casein. *Biochim. Biophys. Acta* 673:487.
 - 220 Saito, T., T. Itoh, S. Adachi, T. Suzuki, and T. Usui. 1981. The chemical structure of neutral and acidic sugar chains obtained from bovine colostrum κ -casein. *Biochim. Biophys. Acta* 678:257.
 - 221 Saito, T., T. Itoh, S. Adachi, T. Suzuki, and T. Usui. 1982. A new hexasaccharide chain isolated from bovine colostrum κ -casein taken at the time of parturition. *Biochim. Biophys. Acta* 719:309.
 - 222 Scatchard, G., J. S. Coleman, and A. L. Shen. 1957. Physical chemistry of protein solutions. II. The binding of some small anions to serum albumin. *J. Am. Chem. Soc.* 79:12.
 - 223 Schmidt, D. G. 1964. Starch-gel electrophoresis of κ -casein. *Biochim. Biophys. Acta* 90:411.
 - 224 Schmidt, D. G., P. Both, and P. J. de Koning. 1966. Fractionation and some properties of κ -casein variants. *J. Dairy Sci.* 49:776.
 - 225 Schmidt, D. G., and T.A.J. Payens. 1963. The purification and some properties of calcium-sensitive α -casein. *Biochim. Biophys. Acta* 78:492.
 - 226 Schmidt, D. G., and T.A.J. Payens. 1976. Micellar aspects of casein. *Surf. Coll. Sci.* 9:165.
 - 227 Schmidt, D. V., and K. E. Ebner. 1971. Isolation and properties of α -lactalbumin from various sources. *Biochim. Biophys. Acta* 243:273.
 - 228 Shahani, K. M., W. J. Harper, R. G. Jensen, R. M. Parry, Jr., and C. A. Zittle. 1973. Enzymes in bovine milk: A review. *J. Dairy Sci.* 56:531.
 - 229 Shimizu, M., C. Kanno, and K. Yamauchi. 1978. Microheterogeneity and some properties of the major glycoprotein fraction isolated from bovine milk fat globule membrane after delipidation. *Agric. Biol. Chem.* 42:981.
 - 230 Singer, S. J. 1974. Molecular organization of membranes. *Annu. Rev. Biochem.* 43:805.
 - 231 Snow, L. D., D. G. Colton, and K. L. Carraway. 1977. Purification and properties of the major sialoglycoprotein of the milk fat globule membrane. *Arch. Biochem. Biophys.* 179:690.
 - 232 Spencer, E. M., and T. P. King. 1971. Isoelectric heterogeneity of bovine plasma albumin. *J. Biol. Chem.* 246:201.
 - 233 Swaisgood, H. E. 1973. The caseins. *CRC Crit. Rev. Food Technol.* 3:275.
 - 234 Swaisgood, H. E. 1975. Primary sequence of kappa-casein. *J. Dairy Sci.* 58:583.
 - 235 Swaisgood, H. E., and J. R. Brunner. 1962. Characterization of κ -casein obtained by fractionation with trichloroacetic acid in a concentrated urea solution. *J. Dairy Sci.* 45:1.
 - 236 Swaisgood, H. E., and J. R. Brunner. 1963. Characteristics of kappa-casein in the presence of various dissociating agents. *Biochem. Biophys. Res. Comm.* 12:148.
 - 237 Swaisgood, H. E., J. R. Brunner, and H. A. Lillevik. 1964. Physical parameters of κ -casein from cow's milk. *Biochemistry* 3:1616.
 - 238 Swaisgood, H. E., B. L. Larson, E. B. Kalan, J. R. Brunner, C. V. Morr, and P.M.T. Hansen. 1975. Methods of gel electrophoresis of milk proteins. *Am. Dairy Sci. Assoc., Champaign, IL.*
 - 239 Swope, F. C., and J. R. Brunner. 1968. The fat globule membrane of cow milk: a reassessment of isolation procedures and mineral composition. *Milchwissenschaft* 23:470.
 - 240 Szuchet-Derechin, S., and P. Johnson. 1965. The "albumin" fraction of bovine milk. I. Overall chromatographic fractionation on DEAE-cellulose. *Eur. Polym. J.* 1:271.
 - 241 Tewari, U. J., and T.K.S. Mukkur. 1975. Isolation and physico-chemical characterization of bovine serum and colostrum immunoglobulin G (IgG) subclasses. *Immunochemistry* 12:925.
 - 242 Thompson, M. P. 1971. α_s - and β -Caseins. Page 117 in *Milk proteins*. Vol. II. H. A. McKenzie, ed. Academic Press, New York, NY.
 - 243 Thompson, M. P., W. G. Gordon, R. T. Boswell, and H. M. Farrell, Jr. 1969. Solubility, solvation, and stabilization of α_s - β -caseins. *J. Dairy Sci.* 52:1166.
 - 244 Thompson, M. P., and C. A. Kiddy. 1964. Genetic polymorphism in caseins of cow's milk. III. Isolation and properties of α_s -caseins, A, B, and C. *J. Dairy Sci.* 47:626.
 - 245 Thompson, M. P., and L. Pepper. 1964. Genetic polymorphism in caseins of cow's milk. IV. Isolation and properties of β -casein A, B, and C. *J. Dairy Sci.* 47:633.
 - 246 Thompson, M. P., N. P. Tarassuk, R. Jenness, H.

- A. Lillevik, U. S. Ashworth, and D. Rose. 1965. Nomenclature of the proteins of cow's milk — second revision. *J. Dairy Sci.* 48:159.
- 247 Tomasi, T. B., Jr., and J. Bienenstock. 1968. Secretory immunoglobulins. *Adv. Immunol.* 9:1.
- 248 Townend, R., T. T. Herskovits, H. E. Swaisgood, and S. N. Timasheff. 1964. The solution properties of β -lactoglobulin C. *J. Biol. Chem.* 239:4196.
- 249 Townend, R., L. Weinberger, and S. N. Timasheff. 1960. Molecular interactions in β -lactoglobulin. IV. The dissociation of β -lactoglobulin below pH 3.5. *J. Am. Chem. Soc.* 82:3175.
- 250 Townend, R., R. J. Winterbottom, and S. N. Timasheff. 1960. Molecular interactions in β -lactoglobulin. II. Ultracentrifugal and electrophoretic studies of the association of β -lactoglobulin below its isoelectric point. *J. Am. Chem. Soc.* 82:3161.
- 251 Trainin, Z., and H. Ungar-Waron. 1981. IgM in the bovine immune system and its relevance to disease. Page 169 in *The ruminant immune system*. J. E. Butler, ed. Plenum Press, New York, NY.
- 252 Trieu-Cuot, P., and J.-C. Gripon. 1981. Electrofocusing and two-dimensional electrophoresis of bovine caseins. *J. Dairy Res.* 48:303.
- 253 Tran, V. D., and B. E. Baker. 1970. Casein. IX. Carbohydrate moiety of κ -casein. *J. Dairy Sci.* 53:1009.
- 254 Vaerman, J. P. 1970. Studies on IgA immunoglobulin in man and animals. Thesis, Sintal-Lonvain, Belgium.
- 255 Vanaman, T. C., K. Brew, and R. L. Hill. 1970. The disulfide bonds of bovine α -lactalbumin. *J. Biol. Chem.* 245:4583.
- 256 Vreeman, H. J., P. Both, J. A. Brinkhuis, and C. Van der Spek. 1977. Purification and some physicochemical properties of bovine κ -casein. *Biochim. Biophys. Acta* 491:93.
- 257 Waite, R., and P. S. Blackburn. 1957. The chemical composition and the cell count of milk. *J. Dairy Res.* 25:328.
- 258 Waugh, D. F., L. K. Creamer, C. W. Slattery, and G. W. Dresdner. 1970. Core polymers of casein micelles. *Biochemistry* 9:786.
- 259 Weber, K., and M. Osborn. 1969. The reliability of molecular weight determinations by dodecyl sulfate-polyacrylamide gel electrophoresis. *J. Biol. Chem.* 244:4406.
- 260 Webster, G. C. 1970. Comparison of direct spectrophotometric methods for the measurement of protein concentration. *Biochim. Biophys. Acta* 207:371.
- 261 Wells, P. W., and P. Eyre. 1971. Preliminary characterization of bovine homocytotropic antibody. *Immunochemistry*. 9:88.
- 262 Wetlaufer, D. B. 1961. Osmometry and general characterization of α -lactalbumin. *C. R. Trav. Lab. Carlsberg.* 32:125.
- 263 Wheelock, J. V., and G. Sinkinson. 1969. Identification of 2-acetamido-2-deoxy-D-galactose in bovine κ -casein. *Biochim. Biophys. Acta* 194:597.
- 264 Wheelock, J. V., and G. Sinkinson. 1973. Carbohydrates of bovine κ -casein. *J. Dairy Res.* 40:413.
- 265 Whitney, R. McL. 1977. Milk proteins. Page 66 in *Food colloids*. H. D. Graham, ed. Avi Publ. Co., Westport, CT.
- 266 Whitney, R. McL. 1983. Milk proteins. In *Fundamentals of Dairy chemistry*. 3rd ed. N. Wong, ed. Avi Publ. Co., Westport, CT (in press).
- 267 Whitney, R. McL., J. R. Brunner, K. E. Ebner, H. M. Farrell, Jr., R. V. Josephson, C. V. Morr, and H. E. Swaisgood. 1976. Nomenclature of the proteins of cow's milk: Fourth revision. *J. Dairy Sci.* 59:795.
- 268 Wilson, A. C., S. S. Carlson, and T. J. White. 1977. Biochemical evolution. *Ann. Rev. Biochem.* 46:573.
- 269 Wooding, F.B.P. 1971. The structure of the milk fat globule membrane. *J. Ultrastruct. Res.* 37:388.
- 270 Woychik, J. H. 1964. Polymorphism in κ -casein of cow's milk. *Biochem. Biophys. Res. Comm.* 16:267.
- 271 Woychik, J. H., E. B. Kalan, and M. E. Noelken. 1966. Chromatographic isolation and partial characterization of reduced κ -casein components. *Biochemistry*. 5:2276.
- 272 Wyckoff, M., D. Rodbard, and A. Chrambach. 1977. Polyacrylamide gel electrophoresis on sodium dodecyl sulfate-containing buffers using multiphasic buffer systems: properties of the stack, valid R_f -measurement, and optimized procedure. *Anal. Biochem.* 78:459.
- 273 Yaguchi, M., D. T. Davies, and Y. K. Kim. 1968. Preparation of κ -casein by gel filtration. *J. Dairy Sci.* 51:473.
- 274 Ziv, G., and S. Gordin. 1973. The composition of bovine udder secretion during the first half of the dry period. *Zentralbl. Veterinärmed. A* 20:285.